

**POLYCHAETOUS ANNELIDS OF THE INDIAN OCEAN INCLUDING
AN ACCOUNT OF SPECIES COLLECTED BY MEMBERS OF
THE INTERNATIONAL INDIAN OCEAN EXPEDITIONS,
1963-1964, AND A CATALOGUE AND BIBLIOGRAPHY
OF THE SPECIES FROM INDIA. PART II***

OLGA HARTMAN

Allan Hancock Foundation, University of Southern California,
Los Angeles, California, U. S. A.

CATALOGUE OF THE POLYCHAETOUS ANNELIDS FROM INDIA

The polychaetous annelids named from India are listed in systematic arrangement, with references to their occurrences in the northern Indian Ocean. The area is here limited by the Red Sea on the west and the western side of the Malay Archipelago on the east. Excluded are many of the species named from the SIBOGA EXPEDITION, 1899-1900, and those from Semper's voyages to the Philippine Islands, named by Grube, 1878.

APHRODITIDAE

Aphrodita aculeata Linnaeus, 1761. Fauvel, 1953, p. 24. Madras.
australis Baird, 1865. Fauvel, 1953, p. 26. Laccadive Sea, in 670 fms.
talpa Quatrefages, 1865. Fauvel, 1953, p. 26. Bay of Bengal.

Aphrogenia alba Kinberg, 1855. Fauvel, 1953, p. 27. Ceylon ; Andaman Islands.

Hermontia bicolor (Grube, 1875 : 67). Red Sea.
erinaceus (Quatrefages, 1865, p. 208). Red Sea.
hystrix (Savigny, 1820). Fauvel, 1953, p. 28. India ; Pillai, 1965, p. 112. Indomalaysia and Red Sea.

Laetmonice filicornis benthaliana McIntosh, 1885. Fauvel, 1953 : 29. Ceylon.

Pontogenia chrysocoma minuta Potts, 1910 : 329. Zanzibar.
indica Grube, 1878. Willey, 1905 : 246, Ceylon ; Fauvel, 1953, p. 29. India and Singapore.
nuda Horst, 1917. Fauvel, 1953, p. 30. Andaman Islands.

* Presented at the 'Symposium on Indian Ocean and Adjacent Seas—Their Origin, Science and Resources' held by the Marine Biological Association of India at Cochin from January 12 to 18, 1971.

The first part of this paper has appeared in Volume 16, No. 1 of the Journal of the Marine Biological Association of India.

POLYNOIDAE

- Admetella longipedata* McIntosh, 1885. Fauvel, 1953, p. 54. Andaman Sea, in 279-569 fms.
- Allmaniella nuchalis*, new species, see Part I.
- ptycholepis* (Grube, 1878). Fauvel, 1953, p. 53. Andaman and Nicobar islands.
- sp. Monro, 1937, p. 260. Arabian Coast, in 1646 m.
- Bouchiria vesiculosa* Wesenberg-Lund, 1949, 4 : 255. Gulf of Iran.
- Drieschia pelagica* Michaelsen, 1892, p. 6. Ceylon. Fauvel, 1953, p. 54. Indian Seas.
- Eunoe pallida* (Ehlers, 1908, p. 49). Indian Ocean, in 616 m ; Fauvel, 1953, p. 39. Andaman Sea and Persian Gulf.
- Gastrolepidia clavigera* Schmarda, 1861, p. 159. Ceylon. Fauvel, 1953, p. 51. Southern India, on nocturnians ; Tampi and Rangarajan, 1964, 6 : 98. Kaapapaia, Ancamans.
- Gattyana deludens* Fauvel, 1932, p. 18. Bay of Bengal. Fauvel, 1953, p. 39. India.
- Harmothoe arabica* Monro, 1937, p. 257. Maldives. Fauvel, 1953, p. 46. Maldives.
- branchiata*, new species, see Part I.
- cornuta* (Potts, 1910, p. 339). Indian Ocean. Monro, 1937, p. 255. Zanzibar, in 786 m.
- dictyophora* (Grube, 1878). Fauvel, 1953, p. 44. Mauras.
- imbricata* (Linnaeus, 1767). Fauvel, 1953, p. 42. Krusadai Island ; Tampi and Rangarajan 1964, 6 : 100. Andaman Islands.
- lunulata* (dele Chiaje, 1841). Monro, 1937, p. 258. South Arabia, in 38 m.
- minuta* (Potts, 1910, p. 337). Indian Ocean. Fauvel, 1953, p. 45. Andaman Islands, in crinoids.
- pseudoampullifera* de Silva, 1965, 30 : 3. Ceylon.
- singawaensis* Izuka, 1912. Fauvel, 1932, p. 23. India ; Fauvel, 1953, p. 48. Madras.
- Herdmanion africanus*, new species, see Part I.
- Herdmanella gracilis* Ehlers, 1908, 16 : 44. Indian Ocean, in 1500-2000 m.
- Hermenia acantholepis* (Grube, 1878). Fauvel, 1953, p. 38. Ceylon.
- Hololepidella commensalis* Willey, 1905, p. 251. Ceylon. Fauvel, 1953, p. 59. Commensal on *Clypeaster*.
- Hyperhalosydna striata* (Kinberg, 1855). Fauvel, 1953, p. 52. Andaman Islands.
- Iphione muricata* (Savigny, 1818). Fauvel, 1953, p. 32. India ; Andaman Islands ; Storch, 1967, 23 : 148. Red Sea.
- Lagisca flaccida* Potts, 1910, 13 : 339. Zanzibar. Fauvel, 1953, p. 41. Ceylon.
- Lepidasthenia grimaldii* (Marenzeller, 1892). Potts, 1910, p. 344. Zanzibar, as *L. maculata*. Fauvel, 1953, p. 56. Arabian Sea.
- microlepis* Potts, 1910. Fauvel, 1953, p. 57. Andaman Islands, on coral stones.
- minikoensis* Potts, 1910, 13 : 344. Indian Ocean.
- ohshimai* Okuda, 1936. Rangarajan, 1963, 5 : 103. Palk Bay, India.
- stylelepis* Willey, 1907, 1 : 260. Persian Gulf.
- Lepidonotus carinulatus* Grube, 1870, 10 : 488. Red Sea. Fauvel, 1953, p. 34. Indian Ocean.
- cristatus* Grube, 1876. Fauvel, 1953, p. 35. India.
- cristatus ornatus* Potts, 1910, 13 : 333. Indian Ocean.
- dictyolepis* Haswell, 1883. Fauvel, 1953, p. 35. Gulf of Mannar.
- fuscirostris* (Schmarda, 1861, p. 152). Ceylon. Fauvel, 1953, p. 38. Ceylon.
- glaucus* (Peters, 1854). Potts, 1910, p. 334. Indian Ocean, as *Lepidonotus carinatus*.
- hedleyi* Benham, 1915. Fauvel, 1953, p. 35. Karachi ; Indian Ocean.
- indicus* Kinberg, 1855, p. 382. Bangka Strait. Fauvel, 1953, p. 47. Ceylon.
- jacksoni* Kinberg, 1855. Fauvel, 1953, p. 34. Ceylon ; Andaman Islands.
- melanogrammus* Haswell, 1883. Fauvel, 1953, p. 36. Andaman Islands.

- purpureus* Potts, 1910, 13 : 334. Zanzibar.
pustulatus Potts, 1910, 13 : 335. Zanzibar.
tenuissetosus (Gravier, 1902, 3 : 222). Red Sea. Fauvel, 1953, p. 36. Orissa and Madras. Tampi and Rangarajan, 1964, 6 : 100. Andaman Islands.
- Paralepidonotus ampullifera* (Grube, 1878). Wesenberg-Lund, 1949, 4 : 253. Iran ; Fauvel, 1953, p. 47. Persian Gulf.
boholensis (Grube, 1878). Fauvel, 1953, p. 47. Persian Gulf.
indicus (Potts, 1910, 13 : 338). Indian Ocean, in 280 fms.
- Polynoe nephrolepidota* Schmarda, 1861, p. 56. Ceylon. Indeterminable.
- Scalissetosus glabrus*, new species, see Part I.
levis Marenzeller, 1902. See Part I.
longicirrus (Schmarda, 1861, p. 336). Ceylon. Fauvel, 1953, p. 50. Ceylon, on echinoderms.
longicirrus Potts, 1910, 13 : 336. Indian Ocean, commensal with gorgoan. Made type of *Australaugeneria pottsi* Pettibone, 1969, 82 : 521.
pellucidus (Ehlers, 1864). Fauvel, 1953, p. 49. South India, on echinoderms.
willei (Potts, 1910, 13 : 340). Indian Ocean.
zeylanica (Wille, 1905), 4 : 250. Ceylon.
- Thormora jukesii* (Baird, 1865). Willey, 1905, 4 : 249. Ceylon, as *Lepidonotus trissochaetus*. Fauvel, 1953, p. 37. Indian Ocean ; Red Sea ; Tampi and Rangarajan, 1964, 6 : 100. Nicobar and Andamans.

POLYODONTIDAE

- Eupanthalis edriophthalma* Potts, 1910, 13 : 345. Indian Ocean. Fauvel, 1953, p. 76. Burma : Ceylon.
- Eupolyodontes sumatranus* Pflugfelder, 1932, p. 282. Sumatra, east coast.
- Panthalis oerstedii* Kinberg, 1855. Fauvel, 1953, p. 78. India.
- Polyodontes flagelliformis* Wesenberg-Lund, 1949, 4 : 261. Gulf of Iran.
maxillosus (Ranzani, 1817). Fauvel, 1953, p. 71. Andaman Sea.
melanonotus Grube, 1878. Fauvel, 1953, p. 72. Indian Seas, in depths to 230 fms.
- Pseudeupanthalis eylathae* Fauvel, 1957, no. 13 : 16. Gulf of Akaba.

PEISIDICIDAE

- Peisidice dorsipapillata* (Marenzeller, 1893). See Part I.

SIGALIONIDAE

- Euthalenessa oculata* (Peters, 1854). Wesenberg-Lund, 1949, p. 258. Iran.
Sthenolepis vulturis Horst, 1917. Monro, 1937, p. 263. Gulf of Aden, in 274 m.
yhleni Malmgren, 1867. Wesenberg-Lund, 1949, 4 : 261. Iran.
- Pholoe* sp., see Part I.
- Psammolyce antipoda* (Schmarda, 1861). Fauvel, 1953, p. 67. Ceylon ; Fauvel, 1957. Gulf of Akaba.
farquharensis Potts, 1910, 13 : 347. Indian Ocean.
fitensis McIntosh, 1885. Fauvel, 1953, p. 67. Mergui Archipelago.
gracilis Potts, 1910, 13 : 348. Indian Ocean.
petersi Kinberg, 1858. Prasad, 1969, 9 : 441. Arabian Sea.
rigida Grube, 1873, 18 : 173. Red Sea. Willey, 1904 : 256. Ceylon.
zeylanica Willey, 1905, 3 : 255. Ceylon. Fauvel, 1953, p. 68. Ceylon.

Sigalion mathildae Audouin and Milne Edwards, 1832. Monro, 1937, p. 262. South Arabia, in 16-22 m.

Sthenelais boa (Johnston, 1839). Fauvel, 1953, p. 61. Ceylon; Indian Ocean.
calcareo Potts, 1910, 13 : 349. Maldives, on atoll; Fauvel, 1953, p. 64. Maldives.
foliosa Potts, 1910, 13 : 351. Seychelles.
laevis Kinberg, 1858. Potts, 1910, 13 : 350. Zanzibar and Maldives, as *S. variabilis glabra*.
longipinnis Grube, 1875, 52 : 77. Red Sea.
orientalis Potts, 1910, 13 : 348. Amirantes, Indian Ocean.
variabilis Potts, 1910, 13 : 349. Maldives and Zanzibar, on atolls; Fauvel, 1953, p. 63. India.
variabilis hirsuta Potts, 1910, 13 : 350. Maldives and Zanzibar; Fauvel, 1953, p. 62. Maldives.
zeylanica Willey, 1905, 4 : 258. Ceylon. Fauvel, 1953, p. 62. Ceylon.

Sthenolepis japonica (McIntosh, 1885). Fauvel, 1953, p. 69. Indian Ocean; Tampi and Rangarajan, 1964, 6 : 101. Andaman Islands, as *Leanira japonica*.

PISIONIDAE

Pisone complexa Alikunhi, 1947, 13 : 105. Madras.
gopalai (Alikunhi, 1941, 13 : 225). Madras.
oerstedii Grube, 1857. Fauvel, 1953, p. 77. Ceylon.
reducta Storch, 1967, 178 : 104. Red Sea.

Pistonidens indica (Aiyar and Alikunhi, 1940, 42 : 106). Madras.

CHRYSOPETALIDAE

Bhawania cryptocephala Gravier, 1902, 3 : 263. Red Sea; Wesenberg-Lund, 1949. Iran.
goodii Webster, 1884. Tampi and Rangarajan, 1964, 6 : 101. Andamans.
myrialepis Schmarda, 1861, p. 164. Ceylon. (may include *B. cryptocephala* Gravier, fide Fauvel, 1953, p. 79).

Chrysopetalum ehlersi Gravier, 1902, 3 : 260. Red Sea. Fauvel, 1953, p. 78, Krusadai Island.

Palmyropsis macintoshi Potts, 1910, 13 : 326. Seychelles; Indian Ocean. Species indeterminate.

AMPHINOMIDAE

Amphinome anatifera Krishnamoorthi and Daniel, 1950, 20 : 17. Madras.
bruguierei Quatrefages, 1865, p. 398. Seychelles.
encopochaeta Schmarda, 1861, p. 143. Ceylon.
indica Schmarda, 1861, p. 142. Ceylon. Indeterminable.
maldivensis Potts, 1909, 12 : 363. Maldives. Questionably *Eurythoe parvecarunculata* Horst.
rostrata (Pallas, 1766). Indian Ocean. Fauvel, 1953, p. 81. Andaman Sea, in 112 fms.

Benthoscolex caecus Horst, 1912, 24 : 38. Ceylon. Fauvel, 1953, p. 93. Ceylon and Laccadive Sea.

Chloeia amphora Horst, 1910, 32 : 172. Malay. Fauvel, 1953, p. 96. Andaman and Nicobar islands.
bengalensis Kinberg, 1867, 24 : 86. India. Indeterminable.
flava (Pallas, 1766). Indian Ocean. Fauvel, 1953, p. 96. Indian Ocean, with bibliography and synonymy.
flava pulchella Baird, 1870. Tampi and Rangarajan, 1964, 6 : 101. Andaman Islands.
flava tumida Baird, 1870, 10 : 232. Indian Ocean. Monro, 1937, p. 254. Maldives, in 27 m.
fusca McIntosh, 1885, 12 : 14. Molucca Islands. Fauvel, 1953, p. 97. Maldives; Fauvel, 1957. Gulf of Akaba.

- longisetosa* Potts, 1909, 12 : 357. Maldives.
maculata Potts, 1909, 12 : 358. Indian Ocean.
parva Baird, 1870, 10 : 233. Fauvel, 1953, p. 96. Ceylon.
rosea Potts, 1909, 12 : 357. Indian Ocean. Fauvel, 1953, p. 97. Persian Gulf.
violacea Horst, 1910, 32 : 174. Malay. Fauvel, 1953, p. 95. Gulf of Oman.
- Eurythoe complanata* (Pallas, 1766). Fauvel, 1951, 23 : 290. Gulf of Akaba. Monro, 1937, p. 252. Red Sea. Tampi and Rangarajan, 1964, 6 : 103. Andamans.
karachiensis Bindra, 1927, 1 : 13. Karachi, Pakistan.
matthai Bindra, 1927, 1 : 12. Indian Ocean. Fauvel, 1953, p. 84. Karachi.
parvecarunculata Horst, 1912. Fauvel, 1953, p. 85. Chilka Lake ; Fauvel, 1953, p. 290. Red Sea ; Tampi and Rangarajan, 1964, 6 : 103. Andamans.
- Hipponea gaudichaudi* Audouin and Milne Edwards, 1833, See Part I.
- Notopygos gardineri* Potts, 1909, 12 : 362. Indian Ocean.
gigas Horst, 1911. Fauvel, 1953, p. 98. Ceylon.
hispidus Potts, 1909, 12 : 359. Indian Ocean. Fauvel, 1953, p. 100. Nicobar Islands, among coral.
hispidus serratus Fauvel, 1917, 56 : 192. Indian Ocean.
labiatus McIntosh, 1885. Fauvel, 1953, p. 99. Indo-Pacific.
variabilis Potts, 1909, 12 : 360. Maldives. Fauvel, 1953, p. 100. Andaman Islands.
- Linopherus acarunculata* (Monro, 1937, 8 : 249). Maldives. Fauvel, 1953, p. 86, as *Pseudeurythoe acarunculata*.
ambigua (Monro, 1933). Fauvel, 1953, p. 90. Maldives, as *Pseudeurythoe*.
hirsutus (Wesenberg-Lund, 1949, 4 : 268). Gulf of Iran, as *Pseudeurythoe hirsuta*.
microcephala (Fauvel, 1932, 12 : 49). Maldives, as *Pseudeurythoe*.
paucibranchiata (Fauvel, 1932, 12 : 47). Gulf of Suez. Fauvel, 1953, p. 86. India.
- Paramphitome indica* Fauvel, 1932, 12 : 51. Arabian Sea. Fauvel, 1953, p. 91. Arabian Sea, in 530 fms.
- Pareurythoe pitipanaensis* de Silva, 1965, 144 : 540. Ceylon.
- Pherecardia polylamellata* de Silva, 1960., Spolia, p. 170. Gulf of Mannar.
striata (Kinberg, 1857). See Part I.

EUPHROSINIDAE

- Euphrosine ceylonica* Michaelsen, 1892, 9 : 1. Ceylon.
foliosa Audouin and Milne Edwards, 1833, 20 : 200. Red Sea. Fauvel, 1953, p. 102. Ceylon ; Nicobars ; Tampi and Rangarajan, 1964, 6 : 103. Andamans.
laureata Savigny, 1818, 3 : 332. Red Sea.
myrtosa Savigny, 1818, 3 : 332. Red Sea. See Fauvel, 1951, 23 : 287, and Fauvel, 1953, p. 101. Ceylon ; Krusadai.
pilosa Horst, 1903. Fauvel, 1951, 23 : 289. Tadjoura. Red Sea.

SPINTHERIDAE

- Spinther mintaceus* Grube, 1860. See Part I.

PHYLLODOCIDAE

- Anatides madetrensis* (Langerhans, 1880). Fauvel, 1953, p. 120. India ; Monro, 1937, p. 265. Red Sea and Arabia.
sancti-josephi (Gravier, 1900), new combination. Red Sea. Willey, 1905, p. 265. Ceylon. See Part I.
- Eteone barantollae* Fauvel, 1932, 12 : 72. Near Calcutta, India.
ornata Grube, 1878. Fauvel, 1953, p. 128. India.
- Eulalia albopicta* Marenzeller, 1879. Fauvel, 1953, p. 123. Nicobar Islands.
manca Gravier, 1900, 2 : 209. Red Sea. (Perhaps *Sige* sp.)
viridis (Linnaeus, 1767). Fauvel, 1953, p. 122. Pamban.

- Eumida sanguinea* (Oersted, 1843). Fauvel, 1953, p. 125. Gulf of Mannar ; Persian Gulf.
- Genetyllis castanea* (Marenzeller, 1879). Willey, 1905, p. 262. Ceylon ; Fauvel, 1953, p. 115. India.
- gracilis* (Kinberg, 1866). Fauvel, 1953, p. 117. Andaman Islands.
- Hesionura elongata* Rao and Ganapati, 1967, p. 10. Indian Ocean, as *Eteonides elongata*.
- serrata* (Hartmann-Schroeder, 1960. 16 : 72). Red Sea.
- ? *Mysta* sp., See Part I.
- Notophyllum splendens* (Schmarda, 1851). Fauvel, 1953, p. 126. Ceylon ; Gulf of Mannar.
- zeylanica* Willey, 1905, pt. 4, p. 262. Mannar Island ; Fauvel, 1953, p. 119. India.
- Phyllodoce dissotyla* Willey, 1905, 4 : 263. Gulf of Mannar. Fauvel, 1953, p. 119. India.
- erythraeensis* Gravier, 1900, 2 : 202. Red Sea.
- foliosopapillata* Willey, 1905, 4 : 264. Ceylon.
- fristedti* Bergstrom, 1914, 3 : 152. Ceylon. Fauvel, 1953, p. 118. Tampian and Rangarajan, 1964, 6 : 104. Malacca Bay.
- gracilis* Gravier, 1900, 2 : 205. Red Sea.
- macrolepidota* Schmarda, 1861, p. 83. Ceylon. Willey, 1905, p. 265. Ceylon.
- malimgreni* Gravier, 1900, 2 : 207. Red Sea. Fauvel, 1953, p. 117. India.
- taprobanaensis* Schmarda, 1861, p. 84. Ceylon. Questionably *Phyllodoce macrolepidota*.
- tenuissima* Grube, 1878. Willey, 1904, p. 265. Ceylon ; Fauvel, 1953, p. 121. Nicobar Islands.
- Pterocirrus ceylonicus* Michaelsen, 1892, 9 : 13. Ceylon. (approaches *Eulalia tenax* Grube)
- Sige macroceros* (Grube, 1860), see Part I.
- Sphaerodoce quadraticeps* (Grube, 1878). Gravier, 1900, 2. Red Sea ; Fauvel, 1953, p. 116. India, as *Phyllodoce*.

ALCIOPIDAE

- Alciopina parasitica* Claparede and Panceri, 1867. Fauvel, 1953, p. 137. India.
- Platohelms capitata* (Greeff, 1876). Fauvel, 1953, p. 138. Ceylon, as *Rhynchonerella fulgens*.
- Torrea candida* (delle Chiaje, 1841). Fauvel, 1953, p. 132. Indian Ocean, as *Asterope candida*.
- Vanadis augeneri* Bonham, 1929. Fauvel, 1951, 23 : 292. Tadjoura.
- formosa* Claparede, 1870. Fauvel, 1953, p. 135. Arabian Sea.

LOPADORRHYNCHIDAE

- Lopadorrhynchus brevis nuchalis* Monro, 1937, 4 : 267. Gulf of Aden.
- (*Prolopadorrhynchus*) *nationalis* Reibisch, 1895. Fauvel, 1951, 23 : 292. Tadjoura, Red Sea.
- uncinatus* Fauvel, 1915. Fauvel, 1953, p. 130. Maldives.
- Pelagobia longicirrata* Greeff, 1879. Fauvel, 1953, p. 131. India.

LACYDONIDAE

- Paralacydonia paradoxa* Fauvel, 1913. See Part I.
- weberi* Horst, 1923. Fauvel, 1953, p. 129. Burma, in 250 fms.

TYPHLOSCOLECIDAE

- Sagittella cornuta* Ehlers, 1913. Fauvel, 1951, 23 : 293. Tadjoura, Red Sea.
- Travelsopsis lobifera* Levinsen, 1885. Fauvel, 1953, p. 139. Arabian Sea.

TOMOPTERIDAE

- Tomopteris* (*Johnstonella*) *aloyst-saubaudiae* Rosa, 1908. Fauvel, 1953, p. 144. Arabian Sea.
(Johnstonella) ducii Rosa, 1908. Fauvel, 1953, p. 143. Bay of Bengal.
(Johnstonella) dunkeri Rosa, 1908, 1 : 276. Ceylon. Fauvel, 1953, p. 145. Red Sea ; Gulf of Siam.
(Johnstonella) mariana Greeff, 1885. Terio, 1953, 20 : 103. Somalia.
(Johnstonella) rofasi Greeff, 1885. Fauvel, 1953, p. 143. Annam ; Gulf of Siam.
(Tomopteris) cavallii Rosa, 1908. Fauvel, 1953, p. 141. North Arabian Sea ; Ceylon.
(Tomopteris) elegans Chun, 1888. Fauvel, 1953, p. 142. Indian Ocean.
(Tomopteris) mortenseni Augener, 1927. Fauvel, 1953, p. 141. Arabian Sea.
(Tomopteris) planktonis Apstein, 1900. Fauvel, 1953, p. 142. Central Arabian Sea.

HESIONIDAE

- Alikunhia elegans* (Alikunhi, 1949, 35 : 192). Madras.
erythraeus (Alikunhi, 1949, 35 : 192). Madras.
longicirrus (Alikunhi, 1949, 35 : 192). Madras.
splendens (Alikunhi, 1949, 35 : 192). Madras.
Gyptis cf arenicolus (La Greca, 1941). See Part I.
? capensis Day, 1963, See above.
lobata (Hessle, 1925). Ceylon.
Hesione genetia Grube, 1866. Fauvel, 1953, p. 105. Ceylon.
intertexta Grube, 1878. Fauvel, 1953, p. 105. Gulf of Mannar ; Tampi and Rangarajan, 1964, 6 : 104. Andamans.
pantherina Risso, 1826. Fauvel, 1953, p. 104. Indian Ocean.
splendida Savigny, 1818, 3. Red Sea. Willey, 1905, p. 266.
Hesionides arenarius Friedrich, 1937. Rao and Ganapati, 1967, p. 56. North Indian Ocean, in beach sand.
gohari Hartmann-Schroeder, 1960, 16 : 74. Red Sea ; Rao and Ganapati, 1967, p. 56. North Indian Ocean.
Leocrates clareddii (Costa in Claparede, 1868). Fauvel, 1953, p. 106. Indian Ocean.
diplognathus Monro, 1926. Fauvel, 1953, p. 107. Mergui Archipelago ; Annam.
Leocratides filamentosus Ehlers, 1908. Fauvel, 1953, p. 107.
Micropathalmus urofimbriata Alikunhi, 1943, 29 : 149. Madras.
Ophiotromus angustifrons (Grube, 1878). Willey, 1905, p. 267. Ceylon, as *Irma limicola*. Fauvel, 1953, p. 109. India.

PILARGIDAE

- Ancistargis brevicirris* Rangarajan, 1964, 6 : 122. Palk Bay, India.
Ancistrosyllis groenlandica McIntosh, 1879. Rullier, 1960, 85 : 236. India.
Ancistrosyllis rigida Fauvel, 1919, p. : 337. Red Sea and India. Fauvel, 1953, p. 110. India.
Loandalia indica Thomas, 1963, 1 : 29. India.
Sigambra constricta (Southern, 1921, p. 573). India. Fauvel, 1953, p. 111. Chilka Lake.
Synelmis albini (Langerhans, 1881), see Part I.

SYLLIDAE

- Autolytus orientalis* Willey, 1905, 4; Ceylon. Fauvel, 1953, p. 162. India.
- Branchiosyllis uncinigera* (Hartmann-Schroeder, 1960, 16 : 86). Red Sea, as *Trypanosyllis uncinigera*.
- Cirrosyllis ceylanica* Schmarda, 1861, p. 78. Ceylon. Indeterminable.
- Eurysyllis tuberculata* Ehlers, 1864, see Part I.
- Eusyllis ceylonica* Augener, 1926, 62 : 453. Fauvel, 1953, p. 159. Ceylon.
homocirrata Hartmann-Schroeder, 1958. Rao and Ganapati, 1967, Northern Indian Ocean.
- Exogone ovalis* Hartmann-Schroeder, 1960, 16 : 106. Red Sea.
remanet Storch, 1966, p. : 182. Red Sea.
simplex Hartmann-Schroeder, 1960, 16 : 107. Red Sea.
- Grubea balani* Hartmann-Schroeder, 1960, 16 : 102. Red Sea.
gracilis Hartmann-Schroeder, 1960, 16 : 101. Red Sea.
nuchalata Hartmann-Schroeder, 1960, 16 : 101. Red Sea.
- Haplosyllis aberrans* Fauvel, 1939, 3 : 290. Indo China.
djiboutiensis Gravier, 1900, 2 : 147. Red Sea.
bisetosa Hartmann-Schroeder, 1960, 16 : 85. Red Sea.
depressa dollfusi Fauvel, 1933, 21 : 50. Gulf of Suez.
spongicola (Grube, 1855). Fauvel, 1953, p. 147. Ceylon; Maldives.
- Langerhansia cornuta* (Rathke, 1843). Fauvel, 1953, p. 153. Gulf of Mannar; Krusadai Island.
- Odontosyllis fulgurans* (Audouin and Milne Edwards, 1833). Fauvel, 1953, p. 160. India.
gibba Claparede, 1863. Fauvel, 1957, Gulf of Akaba.
gibba gravieri Fauvel, 195, 23 : 384. Somaliland.
gracelyi Fauvel, 1928, 34 : 90. Indian Ocean; Fauvel, 1953 : 160. Somaliland.
longicornis Hartmann-Schroeder, 1960, 16 : 95. Red Sea.
- Opisthosyllis brunnea* Langerhans, 1879. Fauvel, 1953, p. 155. Gulf of Mannar; Krusadai Island.
longicirrata Monro, 1939, 41 : 389. Maldives; Red Sea. Fauvel, 1953, p. 154. Maldiv Islands.
papillosa Hartmann-Schroeder, 1960, 16 : 87. Red Sea.
- Paraplonosyllis paucicirra* Hartmann-Schroeder, 1960, 16 : 91. Red Sea.
subterranea Hartmann-Schroeder, 1960, 16 : 92. Red Sea. Rao and Ganapati, 1967, Northern Indian Ocean.
- Parapterosyllis sexoculata* Hartmann-Schroeder, 1960, 16 : 90. Red Sea.
- Parasphaerosyllis indica* Monro, 1937, 4 : 283. Arabia. Fauvel, 1953, p. 162. Arabian Sea.
indica, *fide* Hartmann-Schroeder, 1960, 16 : 84. Red Sea. Homonym.
- Petitla anophthalma* Siewing, 1955. Hartmann-Schroeder, 1960, 16 : 100. Red Sea. Rao and Ganapati, 1967, Indian Ocean.
- Pionosyllis confusa* Hartmann-Schroeder, 1960, 16 : 90. Red Sea.
oculata Hartmann-Schroeder, 1960, 16 : 93. Red Sea. See Part I.
- Pterosyllis formosa* Claparede, 1863. Fauvel, 1951, 23 : 381. Djibouti; Red Sea.
formosa corallicola Hartmann-Schroeder, 1960, 16 : 98. Red Sea.
- Spermosyllis confusa* Hartmann-Schroeder, 1960, 16 : 108. Red Sea.

- Sphaerosyllis brevicirra* Hartmann-Schroeder, 1960, 16 : 105. Red Sea.
capensis serrata Hartmann-Schroeder, 1960, 16 : 103. Red Sea.
minima Hartmann-Schroeder, 1960, 16 : 104. Red Sea. Rao and Ganapati, 1967, Northern Indian Ocean.
xarifae Hartmann-Schroeder, 1960, 16 : 103. Red Sea.
- Streptosyllis reducta* Hartmann-Schroeder, 1960, 16 : 95. Red Sea.
cryptocephala Hartmann-Schroeder, 1960, 16 : 95. Red Sea.
reducta Hartmann-Schroeder, 1960, 16 : 94. Red Sea.
- Syllides papillosa* Hartmann-Schroeder, 1960, 16 : 96. Red Sea.
- Syllis amica* Quatrefages, 1865. Fauvel, 1957. Gulf of Akaba.
curticirris Monro, 1937, 4 : 271. Zanzibar, in 113 m.
gracilis Grube, 1840. Fauvel, 1953, p. 147. Indian Ocean ; Gulf of Mannar ; Maldives.
longissima Gravier, 1900, 2 : 154. Red Sea.
monilaris Savigny, 1818, 3 : 318. Red Sea.
neglecta Grube, 1870, Monatsber., p. 501. Red Sea.
picta Grube, 1870, Monatsber., p. 499. Red Sea.
quadrifasciata Fischli, 1903, 25 : 117. Ternate, Indian Ocean.
violacea Grube, 1870, Monatsber., p. 500. Red Sea.
- Trypanosyllis crosslandi* Potts, 1911, 3 : 14. Zanzibar.
gigantea (McIntosh, 1885). Fauvel, 1953, p. 158. Nicobar Islands.
misakiensis Izuka, 1912. Fauvel, 1953, p. 158. Madras.
taeniaformis (Haswell, 1886), as *Trypanosyllis richardi* Gravier, 1900, 2 : 168. Red Sea.
zebra (Grube, 1860). Fauvel, 1953, p. 157. Indian Ocean ; Persian Gulf.
- Typosyllis bouvieri* Gravier, 1900, 2 : 163. Red Sea.
circummaculata Hartmann-Schroeder, 1960, 16 : 82. Red Sea.
closterobranchia (Schmarda, 1861). Fauvel, 1953, p. 150. Gulf of Mannar ; Andaman Islands.
dentata Hartmann-Schroeder, 1960, 16 : 77. Red Sea.
exilis Gravier, 1900, 2 : 160. Red Sea. Fauvel, 1953, p. 151. Andaman Islands ; Madras ; Maldives.
gerlachei Hartmann-Schroeder, 1960, 16 : 81. Red Sea.
heterocirrata Hartmann-Schroeder, 1960, 16 : 76. Red Sea.
krohnii (Ehlers, 1864). Fauvel, 1953, p. 150. Gulf of Mannar.
lutea Hartmann-Schroeder, 1960, 16 : 81. Red Sea.
magnipectans Storch, 1967, p. 106. Red Sea, as *Syllis* (*Typosyllis*) *magnipectans*.
nuchalis Hartmann-Schroeder, 1960, 16 : 83. Red Sea.
okadai Fauvel, 1934. Fauvel, 1953, p. 152. Andaman Islands.
prolifera (Krohn, 1852). Fauvel, 1953, p. 148. Ceylon ; Gulf of Mannar.
remanei Hartmann-Schroeder, 1960, 16 : 79. Red Sea.
schulzi Hartmann-Schroeder, 1960, 16 : 80. Red Sea.
striata Hartmann-Schroeder, 1960, 16 : 78. Red Sea.
taprobanensis Willey, 1905, pt. 4, p. 268. Gulf of Mannar.
resembles Eusyllis ceylonica, fide Fauvel, 1953, p. 159.
variiegata (Grube, 1860). Fauvel, 1953, p. 148. Ceylon ; Arabian Sea ; Persian Gulf.

NEREIDAE

- Ceratonereis anchylochaeta* (Horst, 1924). Fauvel, 1953, p. 177, as *Nereis*.
burmensis Monro, 1937, 19 : 532. Burma. Fauvel, 1953, p. 196. India.
costae (Grube, 1840). Fauvel, 1953, p. 194. Ceylon.
erythraeensis Fauvel, 1918. Fauvel, 1951, 23 : 251. Somaliland.
flagellipes Fauvel, 1932. Fauvel, 1953, p. 199. India.
hemprichii (Grube, 1873, 51 : 69). Red Sea.
hirchinicola (Eisig, 1870). Fauvel, 1923, 5 : 350. See Part I.
marmorata Horst, 1924. Fauvel, 1951, 23 : 521. Somaliland.
microcephala (Grube, 1878). Fauvel, 1953, p. 198. Siam.
mirabilis Kinberg, 1866. Fauvel, 1953, p. 101. India, as *tentaculata*.
obockensis Gravier, 1902, 3 : 177. Red Sea.
obocki Gravier, 1899, 5 : 240. Red Sea.
pachychaeta Fauvel, 1918. Fauvel, 1953, p. 196. Maldives.

- pectinifera* Grube, 1878. Willey, 1905, p. 272. Ceylon. Referred to *C. costae* by Fauvel, 1953, p. 195. India.
- tripartita* Horst, 1918. Fauvel, 1953, p. 197. Andaman Islands.
- Dendronereides heteropoda* Southern, 1921, 5 : 603. India, in brackish water.
- Dendronereis aestuarina* Southern, 1921, 5 : 598. India, in brackish water.
- arborifera* Peters, 1854. Fauvel, 1953, p. 172. India, in brackish water.
- Leonnates decipiens* Fauvel, 1929, 54 : 180. Gulf of Mannar, India. Fauvel, 1953, p. 171. Gulf of Suez.
- Jousseaumi* Gravier, 1899, 5 : 234. Red Sea ; Fauvel, 1953, p. 171. India ; Suez. Ranganarajan and Sankarankutty, 1961, 3 : 256. On abnormal specimen.
- persica* Wesenberg-Lund, 1949, 4 : 275. Gulf of Aden.
- simplex* Monro, 1939, 41 : 403. North-east Africa.
- Leptonereis davidi* Fishelson and Rullier, 1969, p. 62. Red Sea.
- Lycastopsis cataractarum* Feuerborn, 1932, 5 : 651. Sumatra. Redescribed by Berkeley and Berkeley, 1963, 41 : 907.
- Mastigonereis longicirra* Schmarha, 1861, p. 109. Ceylon. Indeterminable.
- Namalycastis indica* (Southern, 1921). Fauvel, 1953, p. 167. India, in brackish water. de Silva, 1960, p. 173. Florence, 1966, p. 20. Cooum Backwater.
- meraukensis* (Horst, 1918). Fauvel, 1953, p. 166. Mergui, as *Lycastis* ; de Silva, 1960, p. 172.
- meraukensis zeylanica* de Silva, 1961, p. 172. Ceylon.
- nipae* (Pflugfelder, 1933, 105 : 68). Sumatra.
- ranauensis* (Feuerborn, 1932, 5 : 639). Sumatra.
- terrestris* (Pflugfelder, 1933, 105 : 66). Sumatra.
- vivax* (Pflugfelder, 1933, 105 : 69). Sumatra.
- Neanthes augeneri* Gravier and Dantan, 1934. Fauvel, 1951, 23 : 386. Tadjoura, Somaliland.
- belawanensis* (Pflugfelder, 1933, p. 72). Sumatra, as *Nereis*.
- capensis* Willey, 1904. Fauvel, 1953, p. 193. Persian Gulf.
- chilkaensis* (Southern, 1921, 5 : 584). Chilka Lake.
- chingrihattensis* (Fauvel, 1932, 12 : 90). India. Fauvel, 1953, p. 179. India, as *Nereis*.
- criognatha* (Ehlers, 1902). Fauvel, 1953, p. 180. India.
- glisserana* (Horst, 1924). Fauvel, 1953, p. 190. Malay, as *Nereis*.
- glandicincta* (Southern, 1921, 5 : 589). Calcutta. Fauvel, 1953, p. 181. as *Nereis*.
- indica* (Kinberg, 1866). Fauvel, 1953, p. 186. India as *Nereis*.
- longilinguis* Monro, 1937, p. 277. Arabian Sea. Fauvel, 1953, p. 192. Arabian Sea, as *Nereis*.
- meggitti* Monro, 1931, 8 : 580. Rangoon. Fauvel, 1953, p. 194. Rangoon.
- negomboensis* de Silva, 1965, 144 : 543. Ceylon.
- oxypoda* (Marenzeller, 1879). Gravier and Dantan, 1934, 14 : 57. Annam, as *Nereis legeri*.
- reducta* (Southern, 1921, 5 : 593). Chilka Lake. Fauvel, 1953, p. 190. India, as *Nereis*.
- succinea* (Frey and Leuckart, 1847). Pflugfelder, 1933, 72 : 105, Sumatra, as *Nereis belawanensis*.
- unifasciata* (Willey, 1905, 4 : 171). Gulf of Mannar. Fauvel, 1953, p. 182. Gulf of Mannar, as *Nereis*.
- Nereis abnormis* Horst, 1924. Fauvel, 1951, 23 : 387. Tadjoura, Somaliland.
- aegyptia* (Savigny, 1818). Red Sea. Indeterminable ; as *Lycoris*.
- alatsubai* Wesenberg-Lund, 1949, 4 : 281. Gulf of Iran.
- anchylocheta* Horst, 1924. Fauvel, 1953, p. 177. Malacca Strait.
- anomala* (Gravier, 1902, 3 : 191). Red Sea as *Pseudonereis*. Fauvel, 1953, p. 217. India ; Persian Gulf ; Red Sea. Tamoi and Ranganarajan, 1964, 6 : 106. Andamans as *Pseudonereis*.
- cuttacki* Gravier, 1899, 5 : 237. Red Sea. Fauvel, 1953, p. 187. India.
- ehlersiana* Grube, 1878. Willey, 1905, n. 272. Ceylon.
- ehrenbergi* Grube, 1868, 45 : 50. Red Sea.
- ethiopiae* Day, 1957, 38 : 17. Red Sea.

- falcata* (Willey, 1905, 4 : 272). Ceylon, as *Ceratonereis*.
filicaudata Fauvel, 1951, 23 : 519. Somaliland. See Part I.
foliosa Sennarada, 1861, p. 104. Ceylon. Indeterminable.
ghardaqae Hartmann-Schroeder, 1960, 16 : 114. Red Sea.
heteromorpha Horst, 1924. Fauvel, 1953, p. 193. Ceylon.
jacksoni Kinberg, 1865. Fauvel, 1953, p. 169. India.
jacksoni reducta Hartmann-Schroeder, 1960, 16 : 110. Red Sea.
kauderni Fauvel, 1921. Fauvel, 1953, p. 166. India.
onychophora Horst, 1918. Fauvel, 1953, p. 178. India.
persica Fauvel, 1911, as *Nereis zonata persica*. Pruvot, 1930, 70 : 47. Persian Gulf, as *N. longiqua*.
sarsoensis Hartmann-Schroeder, 1960, 16 : 113. Red Sea.
singularis Wesenberg-Lund, 1949, pt. 4, p. 278. Gulf of Iran. Homonym. Resembles *Neanthes oxydopoda* (Marenzeller).
trifasciata Grube, 1878. Fauvel, 1951, 23 : 389. Somaliland ; Fauvel, 1953, p. 183. Maldives.
- Nicon punctata* (Wesenberg-Lund, 1949, 4 : 389). Gulf of Iran. As *Leptonereis*.
- Perinereis affinis* Pruvot, 1930, 70 : 60. Persian Gulf. Resembles *Perinereis striolata* (Grube, 1878).
aiubhitensis Grube, 1878. Fauvel, 1953, p. 209. Andamans. Tampi and Rangarajan, 1964, 6 : 105. Andaman Islands.
brevicirris (Grube, 1869). Tampi and Rangarajan, 1964, 6 : 106. Andaman Islands.
cavifrons Eilers, 1920. Fauvel, 1953, p. 210. Burma ; India.
cultrifera (Grube, 1840). Fauvel, 1953, p. 206. India ; Fauvel, 1957. Gulf of Akaba.
cultrifera brevicirra Grube, 1867. Wesenberg-Lund, 1949, p. 284. Iran.
cultrifera ceylonica de Silva, 1961, 29 : 176. Ceylon.
maindroni Fauvel, 1943, p. 201. Bay of Bengal ; Fauvel, 1953, p. 203. Bengal.
mathaii Aziz, 1938, 1 : 29. Karachi.
nigropunctata (Horst, 1889). Fauvel, 1953, p. 210. India. Includes *Perinereis marmorii* Southern, 1921.
neocaledonia Pruvot, 1930. Fauvel, 1953, p. 211. Arabian Sea ; Tampi and Rangarajan, 1964, 6 : 106. Andamans.
nuntia (Savigny, 1818). Fauvel, 1951, 23 : 525. Somaliland.
nuntia djiboutiensis Fauvel, 1919, p. 420. Djibouti, Red Sea.
nuntia heterodonta Gravier, 1899, 5 : 242. Red Sea.
nuntia valleta (Grube, 1867). Pillai, 1961, 4 : 7. Ceylon.
singaporensis Grube, 1878. Fauvel, 1953, p. 205. Mergui.
suluana Horst, 1924. Fauvel, 1953, p. 204. Andaman Islands.
vancouverica (Ehlers, 1868). Fauvel, 1953, p. 205. Mergui. Includes *P. horsti* Gravier, 1902, 3 : 182. Red Sea.
- Platynereis abnormis* (Horst, 1824). Fauvel, 1953, p. 222. India.
bengalensis Willey, 1905, 4 : 273. Bengal. Fauvel, 1953, p. 218, as *Platynereis dumerilii*, India.
dumerilii (Andouin and M. Edwards, 1833). Fauvel, 1953, p. 218. India.
festiva (Grube, 1874, p. 326). Ceylon.
insolita Gravier, 1902, 3 : 197. Red Sea. See Part I.
pallida Gravier, 1902, 3 : 200. Red Sea.
pulchella Gravier, 1902, 3 : 202. Red Sea. Fauvel, 1953, p. 220. India.
polyscalma Chamberlin, 1919. Fauvel, 1953, p. 221. Andaman Islands.
- Pseudonereis gallapagensis* Kinberg, 1866. Fauvel, 1953, p. 215. Andaman Islands. Tampi and Rangarajan, 1964, 6 : 106. Andaman Islands.
roftnestiana Augener, 1913. Fauvel, 1953, p. 217. Andaman Islands.
- Tambalagamia fauvelii* Pillai, 1961, 4 : 3. Ceylon.
orientalis, new species. See Part I.
- Tylonereis bogoyawlenski* Fauvel, 1911. Fauvel, 1953, p. 168. India. Tampi and Rangarajan, 1964, 6 : 105. Andaman Islands.
fauvelii Southern, 1921, 5 : 582. Chilka Lake. Fauvel, 1953, p. 169. Mergui.

NEPHTYIDAE

- Aglaophamus dibranchis* (Grube, 1877). Fauvel, 1953, p. 225. India; Persian Gulf, as *Nephtys*.
longicephalus, new species, see Part I.
lyratus Kinberg, 1866, 22 : 240. Bangka Strait.
lyrochaetus (Fauvel, 1902). Wesenberg-Lund, 1949, 4 : 293. Iran.
Inermonephtys inermis (Ehlers, 1887). Fauvel, 1953, p. 224. Maldive Archipelago, as *Nephtys*.
Micronephtys sphaerocirrata (Wesenberg-Lund, 1949, No. 4, p. 294. Gulf of Iran, as *Nephtys*.
Nephtys dussumieri Quatrefages, 1865, 1 : 426. Malabar. Questionable.
gravieri Augener, 1913. Fauvel, 1953, p. 226. Bay of Bengal, in 847 fms.
malimgreni Théel, 1879. Fauvel, 1953, p. 226. Andaman Sea, in 279 fms.; Burma, in 250 fms.
oligobranchia Southern, 1921, 5 : 610. India, in brackish water; Fauvel, 1953, p. 228. India.
palatii Gravier, 1904, 10 : 472. Red Sea.
polybranchia Southern, 1921, 5 : 607. Chilka Lake. Fauvel, 1953, p. 227. Madras; Siam.
tulearensis Fauvel, 1919. Wesenberg-Lund, 1949, 4 : 293. Iran.

PHAERODORIDAE

- Clavadorum bengalorum* Fauchald, 1970. Bay of Bengal.

GLYCERIDAE

- Glycera alba cochlinensis* Southern, 1921, 5 : 627. In brackish water. Fauvel, 1953, p. 292. India.
brevicirris Grube, 1870, 47 : 61. Indian Ocean. Questionable.
cinnamomea Grube, 1870. Proc., p. 372. Ceylon.
cirrata, sensu Fauvel, 1932, 12 : 129. Not Grube, 1857. Fauvel, 1953, p. 297. Burma; Madras. Homonym.
convoluta Kieferstein, 1862. Wesenberg-Lund, 1949, 4 : 298. Iran.
embranchiata Krishnamoorthi, 1963, 5 : 97. Madras. Adyar estuary, Madras. (Diagnosis lacking).
gigantea Quatrefages, 1865. Fauvel, 1953, p. 296. Laccadive Sea, in 430 fms.
laccadivae Schmarda, 1861, p. 95. Ceylon. Fauvel, 1953, p. 291. Burma; India; Maldives. As *Glycera edwardsi* Gravier, 1906, 8 : 139. Red Sea.
longipinnis Grube, 1878. Fauvel, 1953, p. 291. Bay of Bengal.
manorae Fauvel, 1932, 12 : 130. Karachi. Fauvel, 1953, p. 298. India.
prashadi Fauvel, 1932, 12 : 126. India. Fauvel, 1953, p. 294. Burma; India; Persian Gulf.
rouxii Audouin and M. Edwards, 1833. Fauvel, 1953, p. 297. India. Pillai, 1961, 4 : 9. Ceylon.
rutilans Grube, in McIntosh, 1885, 12 : 345. Ceylon.
sagittariae Fauvel, 1932, 12 : 127. India. Fauvel, 1953, p. 295. Homonym, perhaps *Glycera tessellata* Grube.
subaenea Grube, 1878. Day, 1967, p. 363. See Part I.
tessellata Grube, 1863. Fauvel, 1953, p. 291. Andaman Islands; Maldives. Fauvel, 1957. Gulf of Akaba.
taprobanensis de Silva, 1965, 144 : 550. Ceylon.

GONIADIDAE

- Glycinde bonhourei* Gravier, 1904, 19 : 474. Red Sea.
oltigodon Southern, 1921, 5 : 629. Chilka Lake. Fauvel, 1953, p. 288. Bay of Bengal, in 840 fms.
- Goniada annulata* Moore, 1905. Fauvel, 1953, p. 283. South of Ceylon, in 660 fms. (Identity may be doubted).
asiatica, new species, see Part I.
emerita Audouin and Milne Edwards, 1833. Fauvel, 1953, p. 282. Cochín. Includes *Goniada longicirrata* Monro, 1937, 4 : 385. Arabian Sea.
eximia Monro, 1937, 4 : 285. North Arabian Sea, in abyssal depth. Homonym.
? maculata Oersted, 1843. Wesenberg-Lund, 1949, 4 : 296. Iran.
multidentata indica Monro, 1937, p. 284. Gulf of Aden; Wesenberg-Lund, 1949, p. 297. Iran.
- Goniadides aciculata* Hartmann-Schroeder, 1960, 16 : 116. Red Sea. Rao and Ganapati, 1967. Northern Indian Ocean.
- Goniadopsis agnesiae* Fauvel, 1928, 34 : 92. Gulf of Mannar. Fauvel, 1953, p. 287. Gulf of Mannar.
incerta Fauvel, 1932, 12 : 122. Burma. Fauvel, 1953, p. 286. Off Burma, in 530 fms.
maskallensis Gravier, 1904, 10 : 475. Red Sea.
- Ophioglycera eximia* (Ehlers, 1900). Fauvel, 1953, p. 285. Fauvel, 1953, p. 285. North Arabian Sea, in 1519-1705 m; as *Goniada eximia*.

ONUPHIDAE

- Diopatra amboinensis* Audouin and Milne Edwards, 1833. Willey, 1905, p. 274. Ceylon.
bengalensis, new species, see Part I.
malabarensis Quatrefages, 1865, p. 346. Malabar. Indeterminable.
neapolitana delle Chiaje, 1841. Fauvel, 1953, p. 252. Burma; India; Maldives; Pillai, 1961, 4 : 13. Ceylon.
neapolitana khargiana Wesenberg-Lund, 1949, 4 : 313. Gulf of Iran.
phyllocirra Schmarda, 1861, p. 133. Ceylon. Indeterminable.
variabilis Southern, 1921, 5 : 611. India, in brackish water. Krishnamoorthi, 1963, 5 : 97. Adyar estuary, Madras.
- Hyalinoecia tubicola* (Müller, 1776). Tampi and Rangarajan, 1964, 6 : 107. Andaman Islands. Fauvel, 1953, p. 261. Bay of Bengal, dredged in 1005 fms.
camiguina Grube, 1878. Willey, 1905, p. 279. Ceylon.
mannarensis Rangarajan and Mahadevan, 1961, 3 : 179. India.
- Nothria conchylega* (Sars, 1835). Willey, 1905, p. 276. Ceylon. Fauvel, 1953, p. 255. Andaman Sea and Ceylon, as *Onuphis*.
holobranchiata (Marenzeller, 1879). Willey, 1905, p. 278. Ceylon. Fauvel, 1953, p. 256. Nicobar Islands; Gulf of Mannar, as *Onuphis*; Wesenberg-Lund, 1949. Iran.
- Onuphis aucklandensis* Augener, 1924. Fauvel, 1953, p. 257. Andaman Islands; off Orissa.
basipicta Willey, 1905, p. 275. Ceylon. Perhaps *Onuphis eremita* Audouin and Milne Edwards.
dibranchiata Willey, 1905, p. 277. Ceylon. Fauvel, 1953, p. 254. Krusadai Island.
eremita Audouin and Milne Edwards, 1833. Fauvel, 1953, p. 257. Burma; Ceylon; Mergui. Krishnamoorthi, 1963, 5 : 97. Adyar estuary, Madras.
furcatoseta Monro, 1937, p. : 290. Red Sea. Fauvel, 1953, p. 254. Gulf of Oman; Gulf of Aden; Red Sea, in 186-375 m.
investigatoris Fauvel, 1932, 12 : 147. Persian Gulf. Fauvel, 1953, p. 258. Arabian Sea, in 35 to 700 fms.
maculata (Baird, 1870, p. 355). Madras.
- Rhamphobranchium bipes* Monro, 1937, p. 298. Gulf of Aden.
chuni Ehlers, 1908. Fauvel, 1953, p. 261. Off India, in 480 and 719 fms.
diversosetosum Monro, 1937, p. 295. Maldives. Fauvel, 1953, p. 262. Maldives, in 183-274 m.

EUNICIDAE

- Eunice afra* Peters, 1854. Fauvel, 1951, 23 : 632. Tadjoura, Somaliland. Fauvel, 1953, p. 235. Mergui. Fauvel, 1957, 13 : 9. Gulf of Akaba. Tampi and Rangarajan, 1964, 6 : 109. Andaman Islands.
- afra paupera* Grube, 1878. Fauvel, 1953, p. 236. India.
- antennata* (Savigny, 1820). Gulf of Suez. Fauvel, 1951, 23 : 633. Tadjoura. Fauvel, 1953, p. 240. Singapore; India; Persian Gulf.
- aphroditidis* (Pallas, 1788). Indian Ocean. Fauvel, 1953, p. 233. India; Maldives.
- australis* Quatrefages, 1865. Fauvel, 1953, p. 240. India.
- coccinea* Grube, 1878. Fauvel, 1953, p. 236. Ceylon; Maldives.
- ehlersi* Gravier, 1900 (4) 2 : 248. Red Sea.
- fauveli* Gravier, 1900 (4) 2 : 236. Red Sea.
- flavo-cuprea* Grube, 1878, p. : 99. Red Sea.
- franklini* Monro, 1924, 36 : 56. Arafura Sea.
- floridana* (Purpura, 1869). Fauvel, 1953, p. 235. Laccadive Sea.
- grubei* Gravier, 1900 (4) 2 : 258. Red Sea. Fauvel, 1951, p. 632. Tadjoura; Fauvel, 1953, p. 237. India.
- guttata* Baird, 1870, 10 : 350. Between Bombay and Singapore.
- indica* Kinberg, 1865. Fauvel, 1953, p. 237. India.
- investigatoris* Fauvel, 1932, 12 : 137. Persian Gulf. Fauvel, 1953, p. 239. Persian Gulf.
- longicirris* Grube, 1870, Monats., p. 492. Red Sea. Fauvel, 1957, p. 6. Gulf of Akaba.
- manorae* Aziz, 1938, 1 : 34. Karachi.
- marenzelleri* Gravier, 1900, (4) 2 : 229. Red Sea. Fauvel, 1953, p. 24. Persian Gulf.
- margaritacea* Fischli, 1903, 25 : 104. Ternate. Questionably *Eunice antennata*.
- (*Nidion*) *gracilis* Crossland, 1904, Proc., p. 327. Zanzibar. Fauvel, 1953, p. 243. India; Mergui.
- pectinata* Grube, 1870, Monats., p. 492. Red Sea.
- perimensis* Gravier, 1900, (4) 2 : 239. Red Sea.
- perrieri* Gravier, 1900, (4) 2 : 232. Red Sea.
- savignyi* Grube, 1878. Fauvel, 1932, 12 : 136. India. Pillai, 1961, 4 : 10. Ceylon; Fauvel, 1953, p. 238. India; Persian Gulf.
- tentaculata* Quatrefages, 1865. Fauvel, 1953, p. 234. India; Laccadive Islands. Tampi and Rangarajan, 1964, 6 : 107. Andaman Islands.
- tubifex* Crossland, 1904, Proc., p. 303. Zanzibar. Fauvel, 1953, p. 232. India; Krusadai Island.
- validobranchiata* Monro, 1937, 8 : 288. South Arabian Coast, in 1046 m.
- Euniphysa aculeata* Wesenberg-Lund, 1949, 4 : 305. Gulf of Iran.
- Lysidice collaris* Grube, 1870, Monats., p. 495. Red Sea. Fauvel, 1953, p. 248. India and Singapore; Fauvel, 1957, 13 : 6. Gulf of Akaba; Tampi and Rangarajan, 1964, 6 : 106. Ternate.
- kuekenhalli* Fischli, 1903, 25 : 106. Ternate.
- Marphysa adenensis* Gravier, 1900, (4) 2 : 270. Red Sea.
- bellii* (Audouin and Milne Edwards, 1833). Wesenberg-Lund, 1949, 4 : 305. Iran.
- borradalei* Pillai, 1958, p. : 94. Tambalagam Lake, Ceylon.
- chevalensis* Willey, 1905, 4 : 282. Gulf of Mannar, India.
- corallina* (Kinberg, 1865). 5 : 239. India.
- graveyi* Southern, 1921, 5 : 617. Chilka Lake. Fauvel, 1953, p. 246. India. Krishna-moorthi, 1963, 5 : 97. Adyar estuary.
- fallax* Marion and Bobretzky, 1875. Fauvel, 1953, p. 247. Gulf of Mannar.
- macintoshi* Crossland, 1903, Proc., p. 137. Zanzibar. Fauvel, 1953, p. 246. Indian Ocean.
- mossambica* (Peters, 1854). Fauvel, 1953, p. 246. India.
- sanguinea* (Montagu, 1815). Crossland, 1903, Proc., p. 141, as *M. furcellata*. Zanzibar. Pillai, 1951, 4 : 12; Tampi and Rangarajan, 1964, 6 : 109. Andaman Islands.
- simplex* Crossland, 1903, Proc., p. 140. Zanzibar. (Homonym of *M. simplex* Langerhans, 1884).
- stragulum* (Grube, 1878). Fauvel, 1953, p. 247. Ceylon.
- teretiuscula* (Schmarda, 1861, p. 129). Ceylon.
- Nemystonereis unicornis* (Grube, 1840). Willey, 1905, 4 : 284. Ceylon; Fauvel, 1953, p. 249. India.

Palola stiliensis (Grube, 1840). Fauvel, 1953, p. 241. India; Fauvel, 1957, 13 : 6. Gulf of Akaba, as *Eunice*. Tampi and Rangarajan, 1964, 6 : 109. Andamans, as *Eunice*.

Paramarphysa orientalis Willey, 1905, 4 : 283. Ceylon. Fauvel, 1953, p. 247. India.

LUMBRINERIDAE

- Lumbrineris bifurcata* (McIntosh, 1885). Wesenberg-Lund, 1949, 4 : 317. Iran.
ebbranchiata (Pallas, 1788, p. 231). Indian Ocean. Indeterminable.
emandibulata Pillai, 1961, 4 : 20. Ceylon.
gracilis (Grube, 1868, 18 : 632). Red Sea.
heteropoda, sensu Fauvel, 1953, p. 268. Bombay; Persian Gulf.
impatiens (Claparède, 1868). Fauvel, 1953, p. 267. India; Maldives; Persian Gulf; Tampi and Rangarajan, 1964, 6 : 110. Andaman Islands.
indica (Kinberg, 1865, p. 569). India. Fauvel, 1953, p. 281. Incompletely known.
inhacae, new species, see Part I.
latreilli Audouin and Milne Edwards, 1834. Fauvel, 1953, p. 266. Ceylon.
mando (Crossland, 1924, Proc., p. 41. Suez.
maxillosa (Ehlers, 1918, 35 : 241). Aru Islands; New Guinea.
nitida (Grube, 1870, Monats., p. 496). Red Sea.
notocirrata Fauvel, 1932, 12 : 156. India. Fauvel, 1953, p. 271. Orissa Coast.
oxychaeta [Gravier, 1900, (4) 2 : 275]. Red Sea.
pallasti Blainville, 1828, 57 : 486. Indian Ocean. Indeterminable.
polydesma (Southern, 1921, 5 : 622). Chilka Lake. Fauvel, 1953, p. 264. Chilka Lake, above high water mark.
pseudobifilaris (Fauvel, 1932, 12 : 154). Burma. Monro, 1937, 8 : 269. Burma, in 250 fms.; off Kerala.
pseudopolydesma Pillai, 1961, 4 : 18. Ceylon.
simplex (Southern, 1921, 5 : 625). India. Fauvel, 1953, p. 264. Chilka Lake.
sphaerocephala (Schmarda, 1861). Fauvel, 1953, p. 267. Andaman Islands; Ceylon. Fauvel, 1957, 13 : 7. Gulf of Akaba.
trigonocephalus (Schmarda, 1861, p. 118). Ceylon. Indeterminable.
Ninoe chilensis Kinberg, 1865. Fauvel, 1953, p. 277. Bay of Bengal, in 105 fms.
pulchra Wesenberg-Lund, 1949, 4 : 319. Gulf of Iran. spp., see Part I.

ARABELLIDAE

- Arabella tricolor* (Montagu, 1804). Fauvel, 1953, p. 274. India. Wesenberg-Lund, 1939, 4 : 318. Iran.
mutans (Chamberlin, 1919). Fauvel, 1953, p. 275. Refers *A. novecrinita* Crossland to *A. mutans*. Tampi and Rangarajan, 1969, 6 : 111. Andaman Islands.
novecrinita logani Crossland, 1924, Proc., p. 75. Suez.
obscura (Willey, 1905) pt. 4, p. 285. Ceylon.
Drilonereis filum (Claparède, 1868). Wesenberg-Lund, 1949, 4 : 318. Iran. Fauvel, 1953, p. 276. Burma. Tampi and Rangarajan, 1964, 6 : 110. Andaman Islands.
major Crossland, 1924, Proc., p. 57. Suez. Fauvel, 1953, p. 277. Bay of Bengal.
monroi new name, see Part I.

LYSARETIDAE

- Oenone fulgida* (Savigny, 1818, 3 : 326). Red Sea. Willey, 1905, 4 : 284, Ceylon. Fauvel, 1953, p. 250. India and Maldives, as *Aglaurides*. Includes *Aglaura erythraeensis* Gravier, 1900, (4) 2 : 278. Suez.

DORVILLEIDAE

- Dorvillea graciloides* Hartmann-Schroeder, 1960, 16 : 117. Red Sea.
gardineri Crossland, 1924, Proc., p. 93. East-Africa. Fauvel, 1953, p. 280. Maldives.
incerta (Schmarda, 1861). Fauvel, 1953, p. 279. Gulf of Mannar.
similis Crossland, 1924, Proc., p. 100. Suez.

ORBINIIDAE

- Naineris laevigata* Grube, 1855. Wesenberg-Lund, 1949, 4 : 323. Iran. Fauvel, 1953, p. 310. Ceylon. Fauvel, 1957, 13 : 7. Gulf of Akaba.
- Orbinia cuvieri persica* (Fauvel, 1932, 12 : 162). Persian Gulf. Fauvel, 1953, p. 301. Persian Gulf.
exarmata (Fauvel, 1932, 12 : 163). Bay of Bengal. Fauvel, 1953, p. 304. Bay of Bengal, in 133 fms.
- Phylo kuppferi* (Ehlers, 1874). Wesenberg-Lund, 1949, 4 : 321. Iran. sp., as *Aricia nuda* Fauvel, 1953, p. 303. Off Burma, in 34 fms. (Not Moore, 1911).
- Scoloplos kerguelensis* McIntosh, 1885. Fauvel, 1953, p. 307. India.
(Leodamas) chevalierii (Fauvel, 1902). Fauvel, 1953, p. 308. Gulf of Mannar. Wesenberg-Lund, 1949, 4 : 322. Iran.
(Leodamas) gracilis Pillai, 1961, 4 : 22. Ceylon.
(Leodamas) latus (Chamberlin, 1919). Fauvel, 1953, p. 309. Off Burma, in 250 fms.
marsupialis Southern, 1921, 5 : 632. Chilka Lake. Fauvel, 1953, p. 306. Krusadai Island. Pillai, 1961, 4 : 22. Ceylon. Tampi and Rangarajan, 1964, 6 : 111. Andamans.

PARAONIDAE

- Aricidea* spp., see Part I.
- Paraonis gracilis* (Tauber, 1879). See Part I.

SPIONIDAE

- Aonides oxycephala* (Sars, 1862). Wesenberg-Lund, 1949, 4 : 324. Iran. Fauvel, 1957, 13 : 7. Gulf of Akaba.
- Laonice brevicristata* Pillai, 1961, 4 : 23. Ceylon.
cirrata (Sars, 1851). Fauvel, 1953, p. 315. Orissa, India.
- Nerine cirratulus* (delle Chiaje, 1828). Fauvel, 1953, p. 312. India.
lefebvrei Gravier, 1905, 11 : 43. Red Sea. Fauvel, 1951, 23 : 635. Tadjoura, Somaliland.
- Nerinides knight-jonesi* de Silva, 1961, 29 : 184. Ceylon. Tampi and Rangarajan, 1964, 6 : 11. Andaman Islands.
williamsi de Silva, 1961, 29 : 183. Ceylon.
- Paraprionospio lamellibranchia*, n. sp., see Part I.
- Polybranchia foxi* Potts, 1928, 22 : 693. Suez Canal.
- Polydora armata* Langerhans, 1880. Watson, 1905, 4 : 325. Ceylon, in sponge : Gulf of Mannar. Fauvel, 1953, p. 321. Ceylon.
ciliata (Johnston, 1838). Fauvel, 1932, 12 : 172. India. Fauvel, 1953, p. 319. Orissa Coast.
coeca (Oersted, 1843). Fauvel, 1953, p. 319. Gulf of Mannar, among sponges.
hornelli Willey, 1905, 4 : 286. Gulf of Mannar. Fauvel, 1953, p. 318. Gulf of Mannar, among oyster shells.
peristomialis, new species, see Part I.

- Prionospio bocki* Söderström, 1920. Wesenberg-Lund, 1949, 4 : 326. Iran.
cirrifer Whén, 1883. Fauvel, 1953, p. 323. India. See Part I.
ehlersi Fauvel, 1928. See Part I.
krusadensis Fauvel, 1929, 54 : 82. Gulf of Mannar.
malmgreni Claparède, 1870. See Part I.
polybranchiata Fauvel, 1929, 54 : 184. Indian Ocean. Fauvel, 1953, p. 324. India.
- Pseudopolydora antennata* (Claparède, 1870). Fauvel, 1953, p. 316. Arabian Sea.
kemp (Southern, 1921, 5 : 636). Calcutta. Fauvel, 1953, p. 317. Salt Lakes, Calcutta.
- Scoleteps indica* Fauvel, 1928, 34 : 93. Gulf of Mannar. Fauvel, 1953, p. 313. Gulf of Mannar; Krusadai Island.
- Spio bengalensis* Willey, 1908, 2 : 389. Bay of Bengal.
multiculata (Rioja, 1919). Rutier, 1963, 87 : 477. Mauritius Island.
- Spiophanes malayensis* Caullery, 1915, 40 : 325. Ceylon, in sponge; Gulf of Mannar.

MAGELONIDAE

- Magelona cornuta* Wesenberg-Lund, 1949, 4 : 328. Gulf of Iran.
obockensis Gravier, 1905, 11 : 44. Red Sea.

DISOMIDAE

- Disoma cirrifer*, new species, see Part I.
orissae Fauvel, 1932, 12 : 174. Indian Ocean. Fauvel, 1953, p. 327. Orissa.
watsoni Fauvel, 1916. Achari, 1968, 19 : 100. Gulf of Mannar, as *Trochochaetus*.

POECILOCHAETIDAE

- Poecilochaetus serpens* Allen, 1904. Achari, 1968, 19 : 102. Pamban, Gulf of Mannar, intertidal.

HETEROSPIONIDAE

- Heterospio longissima* Ehlers, 1875. Arabian Sea. See Part I.

CHAETOPTERIDAE

- Chaetopterus variopedatus* (Renier, 1804). Grube, 1874, Proc., p. 328, Ceylon, as *C. appendiculatus*. Crossland, 1904, Proc., p. 272, 288, as *C. longiramus* and *C. longipes*. Maldives. Gravier, 1906, (4) 8 : 186. Red Sea. as *C. variopedatus djiboutiensis*. Fauvel, 1953, p. 337. India.
- Mesochaetopterus minuta* Potts, 1914, Proc., p. 963. Fauvel, 1953, p. 342. India. Tampi and Rangarajan 1964, 6 : 114. Andaman Islands.
- Phyllochaetopterus aciculiger* Crossland, 1904, Proc., p. 278. Maldives Islands. Fauvel, 1953, p. 341. India.
arabicus Grube, 1870, Monats., p. 507. Red Sea.
elioti Crossland, 1903, Proc., p. 172. Zanzibar. Fauvel, 1953, p. 340. May include *P. gardineri* Crossland, 1904.
herdmani Hornell, in Willey, 1905, 4 : 292. Ceylon.
ramosus Willey, 1905, 4 : 293. Ceylon.
socialis Claparède, 1870. Fauvel, 1953, p. 339. India.
- Telepsavus bonhourei* Gravier, 1905, 11 : 93. Red Sea.

CIRRATULIDAE

Cauleriella typhlops (Willey, 1905, 4 : 295). Gulf of Mannar, as *Heterocirrus typhlops*. Fauvel, 1953, p. 334. India.

Chaetozone sp., see Part I.

Cirratulus auricapillus Grube, 1870, Monats., p. 503. Red Sea. Indeterminable.

cirratus (Müller, 1776). Fauvel, 1953, p. 334. India.

dasylophus (Marenzeller, 1879). Willey, 1905, 4 : 294. Ceylon, as *Cirratulus complanatus*. Fauvel, 1953, p. 333. India.

filiformis Kieferstein, 1862. Wesenberg-Lund, 1949, 4 : 336. Iran. Fauvel, 1953, p. 333. India.

gracilis Grube, 1870, Monats., p. 504. Red Sea. Perhaps identical with *Timarete anchylochaeta*.

nigromaculata Grube, 1870, Monatsber., p. 504. Red Sea. Questionably the same as *Timarete anchylochaeta*.

Cirriformia chrysoderma (Claparède, 1870). Fauvel, 1953, p. 333. India.

filigera (delle Chiaje, 1828). Wesenberg-Lund, 1949, p. 332. Fauvel, 1953, p. 330. India.

Fauvel, 1957, 13 : 5, Gulf of Akaba, as *Audouinia*.

limnocola Kirkegaard and Santhakumaran, 1967, 130 : 213. Bombay Harbour, in burrows in wood.

maryae de Silva, 1961, 29 : 186. Ceylon.

saxatilis Gravier, 1906, (4) : 8 : 154. Red Sea. Resembles *Cirriformia semicincta*.

semicincta (Ehlers, 1905). Fauvel, 1953, p. 330. India.

tentaculata (Montagu, 1808). Day, 1962, 139 : 648. Mombassa.

Dodecaceria fistulicola Ehlers, 1901. Fauvel, 1953, p. 335. India.

joubini Gravier, 1906 (4) 8 : 156. Red Sea.

Tharyx multifilis Moore, 1909. Wesenberg-Lund, 1949, 4 : 332. Iran Fauvel, 1953, p. 334. India. (Doubtfully this species.)

Tharyx spp., see Part I.

Timarete anchylochaeta (Schmarda, 1861). Wesenberg-Lund, 1949, 4 : 331. Iran. Fauvel, 1953, p. 332, as *Audouinia*.

cylindricus (Schmarda, 1861). Willey, 1905, 4 : 294. Ceylon, as *Cirratulus*.

ACROCIRRIDAE

Acrocirrus uchidai Okudai, 1934. See Part I.

Acrocirrus sp., see Part I.

COSSURIDAE

Cossura dayi, new name, see Part I.

Cossurella dimorpha, new genus, new species, see Part I.

Cossurella sp., see Part I.

FLABELLIGERIDAE

Brada mamillata Grube, 1877. Wesenberg-Lund, 1949, 4 : 334. Iran. Fauvel, 1953, p. 352. Arabian Sea, in 555 fms.

talehsapensis Fauvel, 1932, 12 : 184. Gulf of Siam.

sp., papillated. See Part I.

[18]

- Diplocirrus glaucus* (Malmgren, 1867). Fauvel, 1953, p. 353. Mergui, in 5 fms.
- Fauveliopsis arabica*, new species, see Part I.
sp., see Part I.
- Flabelligera diplochaitos* (Otto, 1821). Fauvel, 1953, p. 344. Arabian Sea. Fauvel, 1957, 13 : 8. Gulf of Akaba.
- Flabelligera* sp., see Part I.
- Ilyphagus hirsutus* Monro, 1937, 8 : 304. South Arabian Sea, in 3385 m. Fauvel, 1953, p. 354. (same record).
- Pherusa bengalensis* (Fauvel, 1932, 12 : 180). Bengal and Madras. Fauvel, 1953, p. 347. India.
coronata (Ehlers, 1908, 16 : 121). Indian Ocean, in 141-1362 m.
curviseta (Cautley, 1944). See Part I.
eruca indica (Fauvel, 1928, 34 : 95). Krusadai Island. Fauvel, 1953, p. 347.
flabellata (Sars, 1871). See Part I.
hamocarens (Monro, 1937, 8 : 302). North Arabian Sea, in 759-1024 m. Fauvel, 1953, p. 345. (Same record).
parmata (Grube, 1877). Willey, 1905, 4 : 289. Ceylon. Fauvel, 1953, p. 346. Madras.
plumosa (Müller, 1776). Wesenberg-Lund, 1949, 4 : 334. Iran, as *Stylarioides*.
tenera (Grube, 1877, 18 : 636). Red Sea, as *Siphonostomum*.
- Semiodera bifidus* (Fauvel, 1932, 12 : 182). Arabian Sea. Fauvel, 1953, p. 349. Arabian Sea, as *Stylarioides*.

SCALIBREGMIDAE

- Parasclerocheilus branchiatus* Fauvel, 1928, 34 : 159. Gulf of Mannar. Fauvel, 1953, p. 356. India.
- Scalibregma inflatum* Rathke, 1843. Fauvel, 1953, p. 355. Gulf of Oman, in 609 fms.

OPHELIIDAE

- Ammotrypane aulogaster* Rathke, 1843. Wesenberg-Lund, 1949, 4 : 335. Iran. Fauvel, 1953, p. 359. India.
grandis Pillai, 1961, 4 : 25. Ceylon.
polychetes (Grube, 1866, 43 : 65). Red Sea.
- Armandia lanceolata* Willey, 1905, 4 : 288. Mannar Island. Tampi and Rangarajan, 1964, 6 : 115. Andaman Islands.
leptocirrus Grube, 1878. Wesenberg-Lund, 1949, 4 : 335. Iran. Fauvel, 1957, 13 : 8. Gulf of Akaba. Tampi, 1958, 10 : 15. India.
melanura Gravier, 1905, 11 : 89. Red Sea.
weissenbornii Kuekenenthal, 1887, 21 : 366. Red Sea.
- Polyophthalmus pictus* (Dujardin, 1839). Fauvel, 1953, p. 360. India. Refers *P. ceylonensis* Kuekenenthal, 1887, from Ceylon, *P. collaris* Michaelsen, 1892, from Ceylon, *P. longisetosus* Michaelsen, 1892, from Ceylon, to this species.
- Travisia arborifera* Fauvel, 1932, 12 : 191. Indian Ocean.

STERNASPIDAE

- Sternaspis scutata* (Ranzani, 1807). Wesenberg-Lund, 1949, 4 : 345. Iran.
spp., see Part I.

CAPITELLIDAE

- Barantolla sculpta* Southern, 1921, 5 : 643. Salt Lake, Calcutta. Fauvel, 1953, p. 370. India.
- Branchiocapitella singularis* Fauvel, 1932, 12 : 197. India. Fauvel, 1953, p. 371. India.
- Capitella capitata* (Fabricius, 1780). Stephenson, 1908, 2 : 39. India, as *Matla bengalensis*.
- Capitellethus dispar* Ehlers, 1907. Fauvel, 1953, p. 371. India. Tampi and Rangarajan, 1964, 6 : 115. Andamans.
- Dasybranchus caducus* (Grube, 1846). Wesenberg-Lund, 1949, 4 : 338. Iran. Fauvel, 1953, p. 365. India. Fauvel, 1957, 13 : 8. Gulf of Akaba.
- carneus* Grube, 1870, Monats., p. 505. Red Sea.
- microchaetus* (Schmarda, 1861, p. 56). Ceylon, as *Oncoscolex*. Fauvel, 1953, p. 357. Ceylon.
- Heteromastides platyproctus* Pillai, 1961, 4 : 27. Ceylon, in brackish lake. (Doubtfully this genus.)
- Heteromastus bifidus* Augener, 1914. Fauvel, 1953, p. 358. India.
- deductus* Pillai, 1961, 4 : 29. Ceylon.
- filiformis* (Claparède, 1864). See Part I.
- similis* Southern, 1921, 5 : 640. Chilka Lake. Fauvel, 1953, p. 366. India.
- Leiocapitellides analis* Hartmann-Schroeder, 1960, 16 : 119. Red Sea.
- Leiochrides branchiatus*, new species. See Part I.
- Mastobranchus indicus* Southern, 1921, 5 : 645. Barantolla, India, in brackish lake. Fauvel, 1953, p. 369. India.
- Mediomastus caudatus*, new species. See Part I.
- Notomastus ceylonicus* Pillai, 1961, 4 : 28. Ceylon.
- fauxi* Day, 1955, 42 : 422. India. As *Notomastus giganteus* Fauvel, 1932, 12 : 194. India.
- latericeus* Sars, 1851. Wesenberg-Lund, 1949, 4 : 336. Iran. Fauvel, 1953, p. 364. India.
- zeylanica* Willey, 1905, 4 : 287. Gulf of Mannar. Perhaps *Heteromastides* sp., fide Pillai, 1960, 4 : 29.
- Parheteromastus tenuis* Monro, 1937, 19 : 536. Burma. Fauvel, 1953, p. 369. India. Generic position doubtful, see Part I.
- Pullia armata* Fauvel, 1929, 54 : 184. Gulf of Mannar. Fauvel, 1953, p. 374. Gulf of Mannar.
- Rashgua rubroincta* Wesenberg-Lund, 1949, 4 : 336. Gulf of Iran.
- Scyphoproctus djiboutensis* Gravier, 1904, 10 : 557. Red Sea. Fauvel, 1953, p. 373. India.
- variabilis* Rangarajan, 1963, 5 : 251. Gulf of Mannar.

ARENICOLIDAE

- Arenicola bombayensis* Kewalramani *et al.*, 1959, 11 : 113. Bombay.
- brasiliensis* Nonato, 1958. Tampi and Rangarajan, 1963, 5 : 108. Andaman Islands.
- Tampi and Rangarajan, 1964, 6 : 115. Andaman Islands.

MALDANIDAE

- Asychis gangeticus* Fauvel, 1932, 12 : 206. India.
- Axiiothella australis* Augener, 1914. Tampi and Rangarajan, 1964, 6 : 116. Andaman Islands.
- obockensis* (Gravier, 1905, 11 : 323). Red Sea. as *Axiiothella*.
- tambalagamensis* Pillai, 1961, 4 : 30. Ceylon.

- Clymene amphistoma* Savigny, 1818, 3 : 341. Red Sea. Indeterminable.
diadema Grube, 1868, (4) 2 : 55. Red Sea. Perhaps *Axiotothella* sp.
lunbricalis Savigny, 1818, 3 : 94. Red Sea. Perhaps *Nicomache lunbricalis* (Fabricius, 1780).
uranthus Savigny, 1820, 21. Suez. Indeterminable.
- Euclymene africana* (Gravier, 1905, 11 : 321). Red Sea. as *Clymene*.
annandalei Southern, 1921, 5 : 648. Chilka Lake. Wesenberg-Lund, 1949, p. 340. Iran.
 Fauvel, 1951, 23 : 636. Tadjoura.
insecta (Ehlers, 1905). Krishnamoorthi, 1963, 5 : 95. Adyar estuary, Madras, as *Clymene*.
watsoni (Gravier, 1906), (4) 8 : 198. Red Sea. as *Clymene*.
- Heteromaldane aequalis* Ehlers, 1908, 16 : 133. Off East Africa, in 1362 m.
- Macroclymene monilis* (Fauvel, 1901). Fauvel, 1957, 13 : 7. Gulf of Akaba.
- Maldane glebifex* Grube, 1860. Wesenberg-Lund, 1949, 4 : 342. Iran.
cristata Treadwell, 1923. Wesenberg-Lund, 1949, 4 : 343. Iran. (Identification doubtful).
sarsi tropica Monro, 1937, 4 (8) : 307. Arabian Coast, in 1189-1354 m.
- Maldanella* sp. see Part I.
- Micromaldane bispinosa* Hartmann-Schroeder, 1960, 16 : 121. Red Sea.
- Nicomache interstricta* Ehlers, 1908, 16 : 134. Indian Ocean, in 660 m.
truncata Willey, 1905, 4 : 290. Ceylon.
- Petaloproctus cirratus* Monro, 1937, 4 (no. 8) : 311. South Arabian Coast, in 1046 m.
 sp., see Part I.
- Praxillella gracilis* (Sars, 1861). Wesenberg-Lund, 1949, 4 : 343. Iran.
 sp., see Part I.
- Rhodine gracillior* Tauber, 1879. Wesenberg-Lund, 1949, 4 : 344. Iran. sp., see Part I.

OWENIIDAE

- Myriochele picta* Southern, 1921, 5 : 638. Chilka Lake.
 sp., see Part I.
- Owenia fusiformis* delle Chiaje, 1844. Wesenberg-Lund, 1949, 4 : 345. Iran. Tampi and Rangarajan, 1964, 6 : 116. Andamans.
- orientalis* (Grube, 1878). Willey, 1905, 4 : 290. Ceylon.

SABELLARIIDAE

- Idanthyrus pennatus* (Peters, 1854), Schmarda, 1861. p. 24. Ceylon, as *Hermella bicornis*.
 Tampi and Rangarajan, 1964, 6 : 117. Andaman Islands.
- Lygdamis indicus* Kinberg, 1867, 22 : 350. Bangka Strait.
- Sabellaria alcocki* Gravier, 1906, 12 : 543. Indian Ocean, in 186 m.
chandrae de Silva, 1961, Spolia, p. 188. Ceylon.
pectinata Fauvel, 1928, 34 : 163. Gulf of Mannar.
pectinata intermedia Fauvel, 1932, 12 : 210. Ganges Delta.

PECTINARIIDAE

- Amphictene* sp., see Part I.
- Cistenides aegyptia* (Savigny, 1818, 3 : 90). Gulf of Suez.

- Lagis abbranchiata* Fauvel, 1932, 12 : 215. Cochin.
Pectinaria antipoda (Schmarda, 1861). Tampi and Rangarajan, 1964, 6 : 117. Andamans.
Pectinaria capensis (Pallas, 1778). Linnaeus, 1788, Syst., p. 3752. Indian Ocean, as *Sabella indica*.
nana Wesenberg-Lund, 1949, 4 : 348. Gulf of Oman.
panava Willey, 1905, 4 : 295. Ceylon.
 sp., see Part I.

AMPHARETIDAE

- Amage posterobranchiata* Fauvel, 1932, 12 : 217. Off Ceylon.
 ? *Ampharete* sp., see Part I.
Auchenoplax sp., see Part I.
Irana heterobranchiata Wesenberg-Lund, 1949, 4 : 350. Gulf of Iran.
Lysippe sp., see Part I.
Melinna aberrans Fauvel, 1932, 12 : 221. India. sp., see Part I.
Schistocomus fauveli Hartman, 1955, Endem., p. 42. Madras. ampharetids, see Part I.

TEREBELLIDAE

- Amaeana* sp., see Part I.
Amphitrite rubra (Risso, 1828). Grube, 1870, Monats., p. 509. Red Sea, as *Terebella vinctipes*. Fauvel, 1957, 13 : 9. Gulf of Akaba.
scylla (Savigny, 1820, 21). Red Sea, as *Terebella scylla*.
Eupolymnia labiata (Willey, 1905, 4 : 298). Gulf of Mannar, as *Polymnia*.
nebulosa (Montagu, 1818). Wesenberg-Lund, 1949, 4. Iran; Fauvel, 1957, 13 : 9. Gulf of Akaba, as *Polymnia*.
trigonostoma (Schmarda, 1861). Willey, 1905, 4 : 300. Ceylon, as *Polymnia triplicata*.
triloba (Fischli, 1903, 25 : 122). Ternate, Indian Ocean, as *Lanice*.
Lanice conchilega (Pallas, 1766). Wesenberg-Lund, 1949, 4 : 355. Iran.
socialis (Willey 1905, 4 : 299). Ceylon. as *Polymnia*.
Loimia annulifilis Grube, 1871. Willey, 1905, 4. Ceylon. Wesenberg-Lund, 1949, 4 : 353. Iran, as subsp. of *L. medusa*.
contorta Ehlers, 1908, 16 : 149. Somaliland, in 741 m.
decora Pillai, 1961, 4 : 33. Ceylon.
medusa (Savigny, 1818, 3). Red Sea, as *Terebella*. Krishnamoorthi, 1963, 5 : 97, *variegata* Adyar estuary, Madras.
 (Grube, 1870, Monats., p. 510). Red Sea, as *Terebella*.
Lysilla pambanensis Fauvel, 1928, 34 : 162. Gulf of Mannar.
Nicolea viridis (Schmarda, 1861, p. 41). Ceylon, as *Phyzella*.
Pista fasciata (Grube, 1870, Monats., p. 513). Red Sea, as *Terebella* (*Phyzella*).
herpini Fauvel, 1928, 34 : 160. Gulf of Mannar.
pachybranchiata Fauvel, 1932, 12 : 231. Laccadive Sea.
typha Grube, 1878. Wesenberg-Lund, 1949, 4 : 354. Gulf of Iran. spp., see Part I.
Polycirrus coccineus Grube, 1870, Monats., p. 515. Red Sea. Fauvel, 1951. 23 : 637. Tad-joura. Fauvel, 1957, 13 : 9. Gulf of Akaba.
twisti Potts, 1928, 22 : 698. Suez.
 sp., see Part I.

- Streblosoma cespitosa* (Willey, 1905, 4 : 305). Ceylon, as *Grymaea*. Fauvel, 1957, 13 : 9. Gulf of Akaba.
persica (Fauvel, 1908, 14 : 386). Persian Gulf, as *Grymaea*.
Terebella ehrenbergi Grube, 1870, Monats., p. 511. Red Sea.
inversa (Willey, 1905, 4 : 297). Ceylon, as *Leprea*.
(Phyzelia) atricapilla Grube, 1870, Monats., p. 512. Red Sea. Questionable.
(Phyzelia) ochroleuca Grube, 1870, Monats., p. 512. Red Sea. Questionable.
tilosaula Schmarda, 1861, p. 41. Ceylon. Indeterminable.
Thelepus plagiostoma (Schmarda, 1861). See Part I.
thoracicus (Grube, 1870, Monats., p. 508. Red Sea, as *Terebella*.

TRICHOBRANCHIDAE

- (Aponobranchus perrieri* Gravier, 1905, 11 : 441. Red Sea. Referred to *Terebellides perrieri*).
Terebellides stroemi Sars, 1935. Gravier, 1905, 11 : 441. Red Sea. Wesenberg-Lund, 1949, 4 : 355. Iran.
umbella Grube, 1870, Monats., p. 514. Red Sea. Indeterminable.
Trichobranchus glacialis Malmgren, 1866. See Part I.

SABELLIDAE

- Branchiomma cingulata* (Grube, 1870). Fauvel, 1957, 13 : 10. Gulf of Akaba, as *Sabella*.
 Raja, 1963, 5 : 126. Madras.
 (Includes *Dasychone maculata* Fischli, 1903, 25 : 125. Ternate).
gravelyi (Aziz, 1938, 1 : 46). Karachi, as *Dasychone*.
kumari (Aziz, 1938, 1 : 46). Karachi, as *Dasychone*.
luciosa (Grube, 1870, Monats., p. 517). Red Sea. as *Sabella (Dasychone)*.
Chone sp., see Part I.
Eurato notata (Grube, 1878). Willey, 1905, 4 : 310. Ceylon.
Fabricia acuseta Banse, 1957, 119 : 103. Red Sea.
Fabriciola ghardaqa Banse, 1957, 119 : 103. Red Sea.
Hypsicomus phaeotaenia (Schmarda, 1861, p. 35). Ceylon, as *Sabella* ; Fauvel, 1953, p. 447, with synonymy. Tampi and Rangarajan, 1964, 6 : 119. Andaman Islands.
Jasmineira caducibranchiata Willey, 1905, 4 : 312. Grand Mannar.
Laonome elegans Gravier, 1906, 12 : 41. Red Sea.
indica Southern, 1921, 5 : 662. Chilka Lake.
Megalomma acrophthalmos (Grube, 1878). Willey, 1905, 4 : 306. Ceylon.
claparedi (Gravier, 1906, 12 : 39). Red Sea, as *Branchiomma*.
intermedium (Beddard, 1889, 21 : 261). Mergui Archipelago, as *Branchiomma*.
mushaensis (Gravier, 1906, 12 : 40). Red Sea, as *Branchiomma*. (Perhaps identical with *M. quadrioculatum*).
quadrioculatum (Willey, 1905, 4 : 307). Ceylon, as *Branchiomma*.
Myxicola fauveli Potts, 1928, 22 : 703. Suez.
Potamilla brevithoracica Pillai, 1961, 4 : 35. Ceylon.
ceylonica Augener, 1926, 62 : 470. Ceylon.
ehlersi Gravier, 1906, 12 : 37. Red Sea.
leptochaeta Southern, 1921, 5 : 651. Bengal.

- Sabella fusca* Grube, 1870, Monats., p. 516. Red Sea.
judica Castelnau, 1842, p. 26. Indian Ocean. Indeterminable.
lamyi Gravier, 1906, 12 : 33. Red Sea.
melanochlora Schmarda, 1861, p. 36. Ceylon. Indeterminable.
porifera Grube, 1878. Willey, 1905, 4 : 309. Ceylon. Tampi and Rangarajan, 1964, 6 : 119. Andaman Islands.
Sabellastarte indica (Savigny, 1818, 3). Indian Ocean. Homonym Willey 1905, 4 : 311. Ceylon. Gravier, 1906, 12 : 42. India, as *Eurato sancti josephi*.
indica quinquevalens Willey, 1905, 4 : 311. Ceylon. Perhaps identical with *S. indica*.
Spirographis tricyclia Schmarda, 1861, p. 37. Ceylon. Indeterminable.

SERPULIDAE

- Bounhorella insignis* Gravier, 1905, 11 : 448. Somaliland.
Conopomatus aculconus Pillai, 1960, 3 : 21. Ceylon.
sectoconus Pillai, 1960, 3 : 23. Ceylon.
Crucigera websteri tricornis Gravier, 1908, (4) 10 : 117. Red Sea.
Ditrupa areolina (Müller, 1776). Wesenberg-Lund, 1949, 4 : 358. Iran.
arietina monilifera Fauvel, 1932, 12 : 247. India.
crenata Ehlers, 1908, 13 : 163. Off Zanzibar, in 463 m.
Eupomatus albiceps Grube, 1869, Monats., p. 250. Red Sea. Willey, 1905, 4 : 312. Ceylon.
 Pillai, 1960, 3 : 12, as *Hydroides*.
exaltatus Marenzeller, 1879. Willey, 1905, 4 : 312. Ceylon, as *Hydroides*. Pillai, 1960, 3 : 10. Ceylon, as *Hydroides*.
inornatus (Pillai, 1960, 3 : 12). Ceylon, as *Hydroides*.
uncinatus Philippi, 1844. Treadwell, 1929, no. 392 : 12. Somaliland, as *E. operculata*.
Hydroides bifurcatus Pixell, 1913, 16 : 76. Indian Ocean.
heteroceros Grube, 1868, 18 : 639. Red Sea. Willey, 1905, 4 : 312. Ceylon. Wesenberg-Lund, 1949, 4 : 357. Iran. Pillai, 1960, 3 : 10. Ceylon.
homoceros Pixell, 1913, 16 : 74. Indian Ocean. Wesenberg-Lund, 1949, 4 : 356. Iran.
minax (Grube, 1878). Willey, 1905, 4 : 314. Ceylon. Pillai, 1960, 3 : 8. Ceylon.
monoceros Gravier, 1908 (4) 10 : 115. Red Sea, as *Serpula (Hydroides)*.
multispinosa ternatensis Fischli, 1903, 25 : 130. Ternate. Indian Ocean.
norvegica (Gunnerus, 1768). Fauvel, 1932, 12 : 242. India. Pillai, 1960, 3 : 12. Ceylon.
perezi Fauvel, 1918, 24 : 342. Persian Gulf.
spinosus (Pixell, 1913, 16 : 78). Suez, as *Eupomatus*.
tambalagamensis Pillai, 1961, 4 : 36. Ceylon.
Mercierella enigmatica Fauvel, 1923. Pillai, 1960, 3 : 33. India, distinguishes this from species of *Ficopomatus*.
Neopomatus similis Pillai, 1960, 3 : 32. Ceylon.
similis rugosus Pillai, 1960, 3 : 32. Ceylon.
uschakovi Pillai, 1960, 3 : 32. Ceylon.
Omphalopomopsis fimbriata (delle Chiaje, 1828). See Part I.
langerhanai zeylanica Pillai, 1960, 3 : 25. Ceylon.
Placostegus crystallinus Scacchi in Philippi, 1844. See Part I.
Porosus (Daudin, 1800, p. 45). Indian Ocean, as *Vermetus*.
Pomatoceros caeruleus (Schmarda, 1861). Fauvel, 1957, 13 : 10. Gulf of Akaba.
crucigera Grube, 1862, 39 : 67. Red Sea.
indicus (Daudin, 1800) as *Vermetus indicus*. Indian Ocean.
Pomatoleios crosslandi Pixell, 1913, 16 : 85. Indian Ocean. Fauvel, 1932, 12 : 243. India.
 Pillai, 1960, 3 : 15. Ceylon.

- Pomatostegus polytrema indica* Fauvel, 1928, 34 : 165. Krusadai Island.
stellatus (Abildgaard, 1789). Willey, 1905, 4 : 314. Ceylon, as *P. actinoceros*. Fauvel, 1932, 12 : 246. Pillai, 1960, 3 : 23. Tampi and Rangarajan, 1964, 6 : 120. Andamans.
- Protula bispiralis* (Savigny, 1820) as *Serpula bispiralis*. Red Sea.
nigra-nucha Fiscelli, 1903, 25 : 128, as *Protula (Protulopsis) nigra-nucha*. Ternate, Indian Ocean.
pallata (Willey, 1905, 4 : 316) as *P. (Protulopsis) pallida*. Ceylon. Wesenberg-Lund, 1949, 4 : 361. Iran. Pillai, 1960, 3 : 5. Ceylon. Referred to *P. tubularia* (Montagu) by Fauvel, 1953, p. 471.
- Salmacina dysteri* (Huxley, 1855). Pillai, 1960, 3 : 3. Ceylon.
- Serpula granulosa* Marenzeller, 1884. Willey, 1905, 4 : 316. Ceylon. Pillai, 1960, 3 : 7. Ceylon. Wesenberg-Lund, 1949, 4 : 356. Iran.
vermicularis chevalensis Pillai, 1960, 3 : 7. Ceylon.
- Spirobranchus ceylonensis* Pillai, 1960, 3 : 20. Ceylon.
coutierei (Gravier, 1905, 11 : 445). Red Sea, as *Pomatoceroopsis coutierei*.
gardineri Pixell, 1913. Fauvel, 1967, 13 : 10. Gulf of Akaba.
giganteus (Pallas, 1766). Fauvel, 1951, 23 : 638. Tadjoura. Fauvel, 1957, 13 : 10. Akaba. Tampi and Rangarajan, 1964, 6 : 120. Andaman Islands.
giganteus arabica Monro, 1937, 4 : 317. South Arabia.
giganteus tubinatus Pixell, 1913, 16 : 81. Maie atoll, Indian Ocean.
jousseaumii (Gravier, 1908 (4) 10 : 130). Red Sea, as *Pomatoceroopsis*.
laticapax (Marenzeller, 1884). Pixell, 1913, 16 : 84. Maldives, as *S. maldivensis*. Wesenberg-Lund, 1949, 4 : 358. Iran.
semperi Moersch, 1863, 1 : 405. Willey, 1905, 4 : 318. Ceylon. Wesenberg-Lund, 1949, 4 : 359. Iran. Pillai, 1960, 3 : 17. Ceylon.
semperi acroceros Willey, 1905, 4 : 318. Indian Ocean. Wesenberg-Lund, 1949, 4 : 359. Iran.
semperi papillosus Wesenberg-Lund, 1949, 4 : 361. Iran.
tricornigerus (Grube, 1878). Willey, 1905, 4 : 318. Ceylon. Pillai, 1960, 3 : 2. Ceylon.
- Vermilopsis glandigerus* Gravier, 1908 (4) 10 : 121. Red Sea. Fauvel, 1957, 13 : 10. Akaba.
pygidalis (Willey, 1905, 4 : 318). Ceylon. Pillai, 1960, 3 : 27. Ceylon.
- Vermilia subcrenata* Savigny, 1818, 3 : 370. Indian Ocean. Indeterminable.

SPIROBIDAE

- Eulaeospira orientalis* (Pillai, 1960, 3 : 37). Ceylon. Pillai, 1970, 8 : 138. Ceylon.
- Leodora knight jonesi* (de Silva, 1965, 144 : 559). Ceylon, as *Spirorbis (Laeospira)*. Pillai, 1970, 8. Ceylon.
- Neodexiospira benhami* Pillai, 1970, 8 : 110. Ceylon.
ceylonicus (Pillai, 1960, 3 : 35). Ceylon. Pillai, 1970, 8. Ceylon.
fauveli Pillai, 1970, 8 : 106. Puttalam Lagoon, Ceylon.
karaitivensis (Pillai, 1960, 3 : 37). Ceylon. Pillai, 1970, 8 : 114. Ceylon.
mannarensis Pillai, 1970, 8 : 114. Mannar Lagoon, Ceylon.
tambalagamensis Pillai, 1970, 8 : 118. Tambalagam Lake, Ceylon.
- Pileolaria bernardi* (Caullery and Mesnil, 1897). Indian Ocean. Pillai, 1970, 8 : 138. Ceylon.
gelastnifera Pillai, 1970, 8 : 101. Ceylon.
grandis Pillai, 1970, 8 : 102. Ceylon.
papillatus (Pixell, 1912. Zanzibar. Pillai, 1970, 8 : 138.
superbus (Pillai, 1960, 3 : 37). Ceylon. Pillai, 1970, 8 : 138. Ceylon.
- Spirorbis indicus* Sterzinger, 1909, 118 : 1450. India. HOMONYM.
 (? *Spirillum*) *albus* Moersch, 1863, 1 : 441. Indian Ocean. Indeterminable.
transversus Daudin in Moersch, 1863, 1 : 465. Indian Ocean.
- Stoa ammonitiformis* Serres, 1855, 4 : 239. Red Sea. Indeterminable.
perforans Serres, 1855, 4 : 239. Red Sea. Indeterminable.
spirulaeformis Serres, 1855, 4 : 239. Red Sea. Indeterminable.

ARCHIANNELIDA

Polygordius madrasensis Aiyar and Alikunhi, 1944, 10 : 113. Madras.
urovridis Aiyar and Alikunhi, 1944, 10 : 118. Madras.

Protodrilus indicus Aiyar and Alikunhi, 1944, 10 : 126. Madras.
pierantonii Aiyar and Alikunhi, 1944, 10 : 134. Madras.

Saccocirrus cirratus Aiyar and Alikunhi, 1944, 10 : 134. Madras.
krusadensis Alikunhi, 1948, 14 : 374. Krusadai.
minor Aiyar and Alikunhi, 1944, 10 : 129. Madras.
orientalis Alikunhi, 1946, 15 : 140. Madras.

MYZOSTOMARIA

Myzostomum costatum Leuckart, 1830, Isis, p. 612. Red Sea.
crosslandi Boulenger, 1913, 37 : 102. Red Sea.
cryptopodium Wheeler, 1897, 12 : 248. Calcutta.
gardineri Atkins, 1927, Proc., p. 355. Maldives.
gopalai Subramaniam in George, 1943, 29 : 150. Madras.
insigne Atkins, 1927, Proc., p. 346. Maldives.
rubro-fasciatum von Graff, 1884, 10 : 33. Red Sea.
striata George, 1943, 29 : 150. India.

The total number of families, genera and species is summarised :

ERRANTIA			SEDENTARIA		
Aphroditidae	..	4 genera, 11 spp.	Orbiniidae	..	5 genera, 10 spp.
Polynoidae	..	21 55	Paraonidae	..	2 2
Polyodontidae	..	5 7	Spionidae	..	12 26
Peisidicidae	..	1 1	Magelonidae	..	1 1
Sigalionidae	..	7 22	Disomidae	..	1 3
Pisionidae	..	2 5	Poecilochaetidae	..	1 1
Chrysopetalidae	..	3 5	Heterospionidae	..	1 1
Amphinomidae	..	10 38	Chaetopteridae	..	4 9
Euphrosinidae	..	1 5	Cirratulidae	..	7 21
Spintheridae	..	1 1	Acrocirridae	..	1 2
Phyllodocidae	..	12 27	Cossuridae	..	2 3
Alciopidae	..	4 5	Flabelligeridae	..	8 19
Lopadorhynchidae	..	2 4	Scalibregmidae	..	2 2
Lacydonidae	..	1 2	Opheliidae	..	4 9
Typhloscolecidae	..	2 2	Sternaspidae	..	1 1
Tomopteridae	..	2 9	Capitellidae	..	16 25
Hesionidae	..	8 18	Arenicolidae	..	1 2
Pilargidae	..	5 6	Maldanidae	..	13 29
Syllidae	..	24 78	Oweniidae	..	2 4
Nereidae	..	16 103	Sabellariidae	..	3 6
Nephtyidae	..	4 13	Pectinariidae	..	4 8
Sphaerodoridae	..	1 1	Ampharetidae	..	7 9
Glyceridae	..	1 17	Terebellidae	..	12 33
Goniadidae	..	5 13	Trichobranchidae	..	2 4
Onuphidae	..	5 22	Sabellidae	..	14 30
Eunicidae	..	7 46	Serpulidae	..	19 56
Lumbrineridae	..	2 24	Spirorbidae	..	6 19
Arabiellidae	..	2 7	Archiannelida	..	3 8
Lysaretidae	..	1 1	Myzostomaria	..	1 8
Dorvilleidae	..	1 4			
1281					

The ten largest families in the Indian Ocean fauna are :

Nereidae with 103 species ;	Amphinomidae with 38 ;
Syllidae with 78 ;	Terebellidae with 33 ;
Serpulidae with 56 ;	Sabellidae with 30 ;
Polynoidae with 55 ;	Maldanidae with 29, and
Eunicidae with 46 ;	Spionidae with 26 species

This proportion compares favourably with that for polychaetes coming from world-wide areas. Based on total computed in Hartman (1959) the ten largest families are ranked in the following order :

Polynoidae,	Sabellidae,
Syllidae,	Phyllodocidae,
Nereidae,	Eunicidae,
Serpulidae,	Maldanidae,
Terebellidae,	Spionidae.

The lower prominence of Phyllodocidae in the Fauna for India may be the result of sparser sampling in areas where they are apt to occur.

None of the families named is limited to the Indian Ocean. On the generic level, one notes genera predominantly or wholly in the Indian Ocean. In the Nereidae are such genera as *Tylonereis*, *Tambalagamia*, *Dendronereis*.

Based on present assessments, the total may be expressed thus :

Errantia are named with 30 families, 160 genera and 532 species.
Sedentaria are named with 27 families, 151 genera and 347 species.
Archiannelida are named with 3 genera and 8 species.
Myzostomaria are named with 1 genus and 8 species.
The grand total is 59 families, 315 genera and 883 species.

It can thus be concluded that the polychaete fauna of the Indian Ocean, restricted as indicated above (and not including the fauna of the Malay Archipelago) is the largest of known tropical polychaete fauna.

Five families are newly entered in the fauna for India :

Peisidicidae, with genus *Peisidice*.
Spintheridae, with genus *Spinther*.
Sphaerodoridae, with genus *Clavadorum*.
Paraonidae, with genera *Aricidea* and *Paraonis*.
Heterospionidae, with genus *Heterospio*.
Acrocirridae, with genus *Acrocirrus*.
Cossuridae, with genera *Cossura* and *Cossurella*.

REFERENCES

- ABILDGAARD, P. C. 1789. Beschreibung einer groszen Seeblase, zween Arten des Steinbohrers, einer groszen Sandröhre. *Schr. Ges. naturf. Freunde Berlin*, 9 : 133-146.
- ACHARI, G. P. K. 1968. Studies on new or little known polychaetes from Indian Seas. *Trochochaeta watsoni* (Fauvel) and *Poecilochaetus serpens* J. mar. biol. Ass. India, 19 (1): 99-106.
- AGOARWALA, A. C. 1924. Polychaeta and Gephyrea from Karachi. *Proc. Lahore Philos. Soc.*, 3 : 69-71, 73-75.
- AIYAR, R. G. 1909. The development and breeding habits of a brackish water polychaete worm from Madras, Calcutta. *J. Proc. Asiatic Soc. Bengal*, 15 : 148-149.
- 1924. A preliminary account of some brackish water polychaetes from Madras. *Proc. 11th Indian Sci. Congr.*, p. 112.
- 1930. On the occurrence of young *Polygordius* and *Polygordius* larvae in townet waters in Madras. *Ibid.*, p. 248.
- 1931. An account of the development and breeding habits of a brackish water polychaete worm of the genus *Marphysa*. *J. Linn. Soc. London*, 37 : 398-404.
- 1931. On some polychaete larvae of the Madras Coast. *Proc. 18th Indian Sci. Congr.*, p. 244.
- 1933. On the anatomy of *Marphysa graveyi* Southern. *Rec. Indian Mus.*, 35 : 287-323.
- 1933. Preliminary observations on some polychaete larvae of the Madras Coast and a note on the occurrence in townet water of the larvae of *Chaetogordius* Moore. *J. Madras Univ.*, 5 : 115-150.
- 1933. Hermaphroditism in *Lycastis indica* (Southern). *Curr. Sci., Bangalore*, 3 : 367-368.
- 1939. On the nephridia of *Pelonospio cirrifera* Wirén. *Proc. Acad. Sci., B. Calcutta*, 9 : 57.
- AND K. H. ALIKUNHI 1940. On a new pisionid from the sandy beach, Madras. *Rec. Indian Mus. Calcutta*, 42 : 89-107.
- AND ——— 1940. On the archiannelids occurring in the sandy beach, Madras. *Proc. 27th Indian Sci. Congr.*, p. 151.
- AND ——— 1944. Change of the generic name *Pisionella* Aiyar and Alikunhi, 1940, into *Pisionidens* (Polychaeta). *Curr. Sci. Bangalore*, 12 (4) : 120.
- AND ——— 1944. On some archiannelids of the Madras Coast. *Proc. Nat. Inst. Sci. India*, 10 (1) : 113-140.
- AND N. K. PANIKKAR 1936. The brackish water fauna of Madras. *Proc. 23rd Indian Sci. Congr.*, p. 357.
- AND ——— 1937. Preliminary observations on the swarming habits and lunar periodicity of *Platynereis* sp. from Madras Harbour. *Proc. 24th Indian Sci. Congr.*, pp. 299-300.
- AND ——— 1937. Observations on the swarming habits and lunar periodicity of *Platynereis* sp. from the Madras Harbour. *Proc. Indian Acad. Sci., (B)*, 5 : 245-260.
- AND M. K. SUBRAMANIAM 1936. Hermaphroditism in *Dasychone cingulata* Grube. With list of hermaphroditic polychaetes. *Curr. Sci., Bangalore*, 5 : 240-242.

- ALIKUNHI, K. H. 1941. On the reproductive system of *Praegeria remota* (Southern). *Proc. Indian Sci. Congr.*, 27 : 152.
- 1941. On a new species of *Praegeria* occurring in the sandy beach, Madras. *Proc. Indian Acad. Sci.*, B. 13 (3) : 193-228.
- 1942. On *Praegeria complexa* n.sp. from the sandy beach, Madras. *Proc. Indian Sci. Congr.*, 28 : 173.
- 1943. Note on the occurrence of archiannelids at Krusadai, together with a description of an undescribed species of *Saccocirrus*. *Ibid.*, 29 : 149.
- 1943. On an undescribed hermaphrodite species of *Microphthalmus* (Hesionidae) occurring on the sandy beach at Madras. *Ibid.*, 29 : 149-150.
- 1946. On a new species of *Saccocirrus* from the Madras beach. *Curr. Sci., Bangalore*, 15 : 140.
- 1947. On *Pisone complexa*, n.sp. from the sandy beach, Madras. *Proc. Nat. Inst. Sci. India*, 13 (3) : 105-127.
- 1948. Genital papillae (suckers) and sperm-sacs in the pisionid polychaete. *Pisionidens indica* (Aiyar and Alikunhi). *Proc. 35th Indian Sci. Congr.*, p. 192.
- 1948. Occurrence of polychaetes in fresh water in the river Godavari. *Proc. 36th Indian Sci. Congr.*, p. 159.
- 1948. On a new hermaphrodite species of *Microphthalmus* (Polychaeta, Hesionidae) from the sandy beach, Madras. *J. Roy. Asiatic Soc. Bengal*, 14 (1) : 17-25.
- 1948. On some Archiannelids of the Krusadai Island. *Proc. Nat. Inst. Sci. India*, 14 (8) : 373-383.
- 1949. On *Anophthalmus* (Fam. Hesionidae) a new genus of polychaete with descriptions of four new species from the sandy beach at Madras. *Proc. 35th Indian Sci. Congr.*, p. 192.
- 1951. On the reproductive organs of *Pisone remota* (Southern), together with a review of the family Pisionidae (Polychaeta). *Proc. Indian Acad. Sci., Sec. B*, 33 (1) : 14-31.
- ANDREW, W. N. 1953. Some annelid and sipunculid worms of the Bimini region. *Amer. Mus. Nov.*, 1617, pp. 1-16.
- ANNANDALE, N. 1907. The fauna of brackish ponds at Port Canning, Lower Bengal. *Rec. Indian Mus., Calcutta*, 1 : 35-74.
- ATKINS, D. 1927. Report on the Myzostomida collected in Torres Strait. *Proc. Zool. Soc., London*, pt. 2, pp. 339-357.
- AUGENER, H. 1927. Die Polychaeten der Sammlung Thilenius von Neuseeland und Samoa. *Mitt. Zool. Mus. Berlin*, 13 (2) : 337-363.
- AZIZ, N. D. 1932. The ecology of polychaetes of Karachi. *Proc. Indian Sci. Congr. Calcutta*, 19 : 261.
- 1938. Fauna of Karachi. Polychaetes. *Mem. Dept. Zool. Punjab Univ.*, 1 : 19-57.
- 1950. Polychaetes of the Arabian Sea. *Proc. 2nd Pakistan Sci. Conference, Dacca*, 2 : 79.
- BAIRD, W. 1865. On new tubicolous annelids in the collections of the British Museum. *J. Linn. Soc. Zool. London*, 8 : 157-160.
- BAL, C. 1933. Note on the occurrence and external features of a fresh water polychaete. *Proc. 20th Indian Sci. Congr.*, pp. 261-262.

- BANSE, KARL. 1957. Die Gattungen *Oriopsis*, *Desdemonia* und *Augeneriella* (Sabellidae, Polychaeta). *Vidensk. Medd. Dansk naturh. Foren.*, 119 : 67-105.
- . 1959. On marine Polychaeta from Mandapam (South India). *J. mar. biol. Ass. India*, 1 (2) : 165-177.
- BERKELEY, E. AND C. BERKELEY. 1963. The proboscis of *Lycastopsis catarractarum* Feuerborn. *Canad. J. Zool.*, 41 : 907-908.
- BINDRA, S. S. 1924. Note on the anatomy of a probable new species of polychaete worm from Karachi. *Proc. 10th Indian Sci. Congr.*, p. 163.
- BORRADAILE, L. A. 1902. On the spawn and young of a polychaete worm of the genus *Marphysa*. *Proc. Zool. Soc. Lond.*, pt. 2, pp. 714-720.
- BOULENGER, C. L. 1913. Report on the Myzostomidae collected by Mr. Cyril Crossland in the Red Sea in 1905. *Proc. Zool. Soc. Lond.*, pp. 85-108.
- CASTELNAU, F. L. 1842. L'Histoire Naturelle des Annelides. Lucas, P. H. (Ed.) In: *Histoire Naturelle des Crustacés, des Arachnides et des Myriapodes*. Paris, pp. 1-46.
- CHAKRAVORTY, M. 1937. On *Paravorticella lycastis* n.sp., an ectoparasitic ciliate on the parapodia of an Indian polychaete, *lycastis indica*. *J. Roy. Micr. Soc. Lond.*, 57 (3) : 71-79.
- CHERIAN, P. V. 1966. Polychaetes from the Cochin Harbour area. *Bull. Dept. Mar. Biol. and Oceanogr. Univ., Kerala*, 2 : 41-50.
- DANIEL, A. AND A. GHOSH. 1964. On some polychaetes from Little Andamans. *Curr. Sci., India*, 33 (8) : 249.
- DAUDIN, F. M. 1800. Nouveau genre de ver a tube calcaire, voisin des Serpules et des dentales Vaginelle. *Bull. Soc. Philom. Paris*, 2 (43) : 145.
- DAY, J. H. 1957. The Polychaete Fauna of South Africa, New species and records from Natal and Mosambique. *Ann. Nat. Mus.*, 14 : 59-129.
- . 1962. Polychaeta from several localities in the western Indian Ocean. *Proc. Zool. Soc. Lond.*, 139 (4) : 627-656.
- . 1965. Some Polychaeta from the Israel South Red Sea Expedition 1962. *Israel South Red Sea Exped., Rep.*, 7 : 15-27.
- . 1967. Polychaeta of Southern Africa. Pt. 1. Errantia ; Pt. 2. Sedentaria. Published by the British Museum of Natural History, pp. 1-878.
- EHLERS, E. 1908. Wissenschaftliche Ergebnisse der deutschen Tiefsee-Expedition auf dem Dampfer *Valdivia* 1898-1899. *Die Bodensässigen Anneliden aus dem Sammlungen der deutschen Tiefsee-Expedition*, 16 (1) : 1-168.
- FAUCHALD, K. 1970. Sphaerodoridae (Polychaeta, Errantia) from world-wide areas. *Los Angeles Co. Mus. Nat. Hist., Bull.* (In press).
- FAUVEL, P. 1908. Sur un Terebellien nouveau de golfe Persique (*Grymaea persica*, n. sp.). *Bull. Mus. Hist. nat. Paris*, 14 (7) : 386-388.
- . 1918. Annélides polychètes des cotes d'Arabie recoltées par M. C. Pères. *Ibid.*, 24 (5) : 329-344.
- . 1919. Annélides polychètes nouvelles de l'Afrique Orientale. *Ibid.*, 25 : 33-39.
- . 1919. Annélides polychètes des Iles Gambier et Touamotou. *Ibid.*, 25 (5) : 336-343.
- . 1934. Indian polychaetes. *Nature, London*, 133 : 143.
- . 1938. Analogies des Annélides polychètes pelagiques de l'Océan Indien et de l'Atlantique. *C. R. Soc. Biogeogr.*, 15 (126) : 30-31.

- 1951. Annélides polychètes du Golfe de Tadjoura (Somaliland) recueillies par M. J. L. Dantan en 19er, au cours de peches nocturnes a la lumière. *Bull. Mus. Hist. nat. Paris*, 23 : 287-294 ; 381-387 ; 519-526 ; 630-640.
- 1953. The Fauna of India including Pakistan, Ceylon, Burma and Malaya. Annelida Polychaeta. *The Indian Press, Allahabad*, 507 pp.
- 1955. Annélides polychètes de la Croisière de la *Calypso* en Mer Rouge en 1952. Results Scientifiques des Campagnes de la *Calypso*. *Campagne en Mer Rouge (1951-1952)*. Masson et Cie, Editeurs, pp. 101-120.
- 1955. Contribution a la faune des Annélides Polychaètes des Cotes d'Israel. State of Israel Ministry of Agriculture, Division of Fisheries. *Sea Fish. Res. Sta., Bull.*, 10, 12 pp. Sphaerodoridae (Polychaeta, Errantia) from world-wide areas.
- 1957. Contributions to the knowledge of the Red Sea. *Los Angeles Co. Mus. Nat. Hist., Bull.* (In Press). Sur Quelques Annélides Polychètes du Golfe d'Akaba. *Fish. Res. Sta. Haifa, Bull.*, 13 : 3-11.
- 1957. Contributions to the knowledge of the Red Sea. Sur quelques Annélides polychètes du Golfe d'Akaba. *Bull. Sea. Fish. Res. Sta.*, 16 : 1-8.
- 1957. Contribution a la faune des Annélides polychètes des Cotes d'Israel. *Bull. Res. Council Israel*, 6B : 213-219.
- FISHELSON, L. AND F. RULLIER 1969. The Second South Red Sea Expedition, 1965. *Rep. No. 9. Quelques Annélides polychètes de la Mer Rouge. Israel J. Zool.*, 18 : 49-117.
- FISCHLI, H. 1903. Polychaeten von Ternate. *Abh. Senckenberg. Naturf. Ges.*, 25 : 90-136.
- FLORENCE, M. R. 1966. Studies on weight changes in *Lycastis indica* Southern. *J. mar. biol. Ass. India*, 8 (1) : 20-27.
- GALLARDO, V. A. 1968. Polychaeta from the Bay of Nha Trang, South Viet Nam. *Naga Rep. Scripps Inst. Oceanogr.*, 4 (3) : 35-227.
- GANAPATI, P. N. AND Y. RADHAKRISHNA 1958. Studies on the polychaete larvae in the plankton off Waltair Coast. *Andhra Univ. Mem. Oceanogr.*, 2 : 210-237.
- M. Y. LAKSHMANA RAO AND R. NAGABHUSHANAM 1958. Biology of fouling in the Visakhapatnam Harbour. *Ibid.*, 2 : 193-209.
- GEORGE, A. I. 1943. On an undescribed species of *Myzostoma* from the Madras harbour, with a note on the occurrence of *Myzostoma adhaerens*. *Proc. Indian Sci. Congr.*, 29 (3) : 150.
- 1950. On *Myzostoma striata* n.sp. from the Madras harbour. *Proc. Indian Acad. Sci., Sec. B.*, 32 : 215-221.
- GHOSH, A. 1963. On a collection of Polychaeta from the south-east coast of India with a new eunicid record. *J. mar. Biol. Ass. India*, 5 : 239-245.
- GIDEON, P. W., P. K. B. MENON, S. R. V. RAO AND K. V. JOSE 1957. On the marine fauna of the Gulf of Kutch. A preliminary survey. *J. Bombay Nat. Hist. Soc.*, 54 : 690-706.
- GRAFF, L. VON 1884. Report on the Myzostomida collected during the voyage of H.M. *Challenger* during the years 1873-76. *Rep. Challenger*, 10 : 1-82.
- 1877. Report on the Myzostomida. *Rep. Challenger, Suppl., Ed.*, 20 : 1-16.
- GRAVIER, C. 1899-1906. Contribution a l'étude des Annélides polychètes de la Mer Rouge. *Bull. Mus. Hist. nat. Paris*, 5 : 234-444 (1899). Nephtyidiens, Glyceriens, 10 : 472-476 (1904) ; Sur un type nouveau de la famille des Capitelliens, *Scyphoproctus* nov. gen., *djiboutensis* n.sp. 10 : 557-561 (1904) ; Cirratuliens, Spionidiens, Ariciens, 11 : 42-46 (1905) ; Flabelligeriens, Opheliens, Capitelliens, Chetopteriens, 11 : 89-94 (1905) ; Cirratuliens, Maldaniens, Amphicteniens, Terebelliens, 11 : 319-326 (1905), 451-454 (1905) ; Sabelliens, 12 : 33-43 (1906) ; Serpuliens, 12 : 110-115 (1906).

- 1906. Sur les affinités de la faune annélide de la Mer Rouge. *Ibid.*, 12 : 149-154.
- 1906. Sur la faune annélide de la Mer Rouge et ses affinités. *C. R. Acad. Sci. Paris*, 142 : 410-412.
- GRUBE, A. E. 1855. Beschreibung neuer oder wenig bekannter Anneliden. *Arch. Nat. Berlin*, 21 (1) : 81-128.
- 1862. Mittheilungen über die Serpulen, mit besonderer Berücksichtigung ihrer Deckel. *Jbr. Schles. Gesells. Breslau*, 39 : 53-69.
- 1866. Resultate einer Revision der Euniceen. *Ibid.*, 44 : 66-68.
- 1870. Mehrere neue Euniceen von die Schiffer-und Fidsche Inseln. *Ibid.*, 47 : 55-56.
- 1875. Bemerkungen ueber die Familie der Aphroditeen. *Ibid.*, 52 : 57-79.
- 1876. Bemerkungen ueber die Familie der Aphroditeen. *Ibid.*, 53 : 46-72.
- 1881. Beschreibungen von neuen Anneliden des zoologischen Museums zu Berlin. *Sitzber. Gesells. nat. Freunde Berlin*, pp. 109-117.
- HARTMAN, O. 1955. Endemism in the North Pacific Ocean, with emphasis on the distribution of marine annelids, and descriptions of new or little known species. *Essays in the Natural Sciences, Los Angeles, California*, pp. 39-60.
- 1959. Catalogue of the Polychaetous Annelids of the World. *Allan Hancock Fdn. Publ. Occas. Pap.*, 23 : 1-628.
- 1965. Supplement and Index to the Catalogue of the Polychaetous Annelids of the World. *Ibid.*, 1-197.
- 1970. Polychaetous Annelids in the Mozambique Channel collected by the International Indian Ocean Expeditions. (in preparation).
- HARTMANN-SCHROEDER, G. 1960. *Polychaeten aus dem Roten Meer. Kieler Meeresforsch.*, 16 : 69-125.
- HASWELL, W. A. 1878. On six new species of annelids belonging to the family Amphinomidae. *Proc. Linn. Soc. New South Wales*, 3 : 341-347.
- HERDMANN, W. A. 1904. Phosphorescence phenomenon (Heteronereis) in the Indian Ocean. *Brit. Ass. Adv. Sci. 73rd Meet.*, pp. 695-696.
- HESSLE, C. 1925. Einiges ueber die Hesioniden und die Stellung der Gattung *Ancistrosyllis*. *Ark. Zool. Stockholm*, 17 (10) : 1-36.
- HORNELL, J. 1903. Report on the pearl-oyster fisheries in the Gulf of Mannar. *Madras Fish. Bull.*, 8 : 23-41.
- HORST, R. 1903. New species of the genus *Euphrosyne* from the Siboga-Expedition, with a Table of the species hitherto known. *Notes Leyden Mus. Jentink*, 23 : 213-222.
- 1910. On the genus *Chloea* with some new species from the Malay Archipelago, partly collected by the Siboga-Expedition. *Ibid.*, 32 : 169-175.
- 1916. A contribution to our knowledge of the SIGALIONINAE. *Zool. Meded. leyden*, 2 : 11-14.
- JOHNSON, H. P. 1903. Freshwater nereids from the Pacific Coast and Hawaii, with remarks on freshwater Polychaeta in general. *Mark Anniv. Vol.*, N.Y., Henry Holt & Co., pp. 205-222.
- KESWAL, M. 1892. A tubicolar annelide. *J. Bombay Nat. Hist. Soc. India*, 7 : 114-115.
- KEWALRAMANI, H. G., P. V. WAGH AND M. R. RANADE, 1959. Taxonomy of the lugworm found off Bombay. *J. Zool. India*, 11 (2) : 109-115.

- KINBERG, J. G. H. 1855. Nya slægten och arter af Annelider. *Oefv. Vet. Akad. Stockholm, Forh.*, 12 : 381-388.
- KIRKEGAARD, J. B. AND L. N. SANTHAKUMARAN 1967. On a new species of annelid associate of marine wood borers : *Cirriformia limnocola* n.sp. (Polychaeta). *Vidensk. Medd. Dansk Naturhist. Forening*, 130 : 213-216.
- KRISHNAMOORTHY, B. 1951. Studies on the osmotic properties of the eggs and larvae of a brackish water polychaete, *Marphysa gravelyi* Southern. *Proc. Indian Acad. Sci.*, 34 (B) : 199-209.
- 1962. Salinity tolerance and volume regulation in four species of polychaetes. *Ibid.*, 56 : 363-371.
- 1963. Gross morphology and histology of nephridia in four species of polychaetes. *Ibid.*, 57 : 195-209.
- 1963. On the distribution of six species of polychaetes in the Adyar estuary, Madras. *J. mar. biol. Ass. India*, 5 : 97-102.
- 1963. Chloride regulation in *Marphysa gravelyi* Southern. *Curr. Sci., Bangalore*, 32 : 463-464.
- 1963. Volume regulation in eggs, larvae and adults of a brackishwater polychaete, *Diopatra variabilis* (Southern). *Proc. Indian Acad. Sci.*, 57 : 275-289.
- 1967. Physiological studies on *Marphysa gravelyi* Southern. Tissue respiration. *Helgoländ Wiss. Meeresunters.*, 16 (1-2) : 179-187.
- AND A. DANIEL 1950. A rare case of a commensal in a polychaete, *Amphinome anatifera* n.sp. *J. Madras Univ., Ser. B*, 20 : 17-24.
- AND S. KRISHNASWAMY 1963. Activity of *Marphysa gravelyi* Southern (Polychaeta) under heterosmotic conditions. *Proc. Indian Acad. Sci.*, 57 (B. 2) : 83-87.
- AND ——— 1966. Physiological studies on *Marphysa gravelyi* Southern. *Ibid.*, 63 (1) : 26-34.
- AND ——— 1966. Physiological studies on *Marphysa gravelyi* Southern. Regulation of body fluid concentration. *Zool. Jahrb. Physiol.*, 72 : 316-326.
- AND ——— 1968. Physiological studies on *Marphysa gravelyi* Southern. Structure of nephridia. *J. mar. biol. Ass. India*, 8 (2) : 244-252.
- KRISHNAN, G. 1936. The development of *Diopatra variabilis* (Southern). *Zeits. wiss. Zool. Leipzig*, 147 : 513-525.
- 1946. Studies on polychaetes from Madras. Thesis. *Madras Univ., ser B*.
- 1950. On the nephridia and coelomoducts of Serpulimorpha and Cirratulidae. *Proc. Nat. Inst. Sci. Calcutta*, 16 : 29-39.
- KURIYAN, G. K. 1950. The fouling organisms of pearl-oyster cages. *J. Bombay Nat. Hist. Soc.*, 49 : 90-92.
- LA GRECA, M. 1946. Studi sui Policheti del Golfo di Napoli. *Pubbl. Staz. Zool. Napoli*, 20 : 270-280.
- MCINTOSH, W. C. 1924. Preliminary note on a collection of South African Polychaetes. *Rep. Union S. Africa Fish. Mar. Biol. Survey*, no. 3, p. 1 ; no. 5, pp. 1-93.
- MARENZELLER, E. 1902. Suedjapanische Anneliden. Aphroditea, Eunicea. *Denkschr. Akad. Wiss. Vienna*, 72 : 563-582.
- MOERCH, O. A. L. 1863. Revisio critica Serpulidarum. Et bidrag til røroremes Naturhistorie. *Naturh. Tidsskr. H. Kroyer, Kobenhavn*, 1 (3) : pp. 347-470.
- MUKUDA, S. 1934. The polychaete genus *Acrocirrus* from Japanese waters. *J. Fac. Sci. Hōkkaidō Imp. Univ.*, 3 (6) : 169-175.

- PALLAS, P. S. 1766. *Miscellanea Zoologica. Hagae Comitum*, pp. 1-224.
- PANIKKAR, N. K. AND R. GOPALA AIYAR. 1937. The brackish water fauna of Madras. *Proc. Indian Acad. Sci.* (6) Sec. B, no. 5, pp. 284-337.
- AND ———. 1939. Observations on breeding in brackish water animals of Madras. *Proc. Indian Acad. Sci.*, Sec. B. 9 (6) : 342-364.
- PAUL, M. D. 1937. Sexual maturity of some sedentary organisms in the Madras harbour. *Curr. Sci., Bangalore*, 5 (9) : 478-479.
- . 1942. Studies on the growth and breeding of certain sedentary organisms in the Madras harbour. *Proc. Indian Acad. Sci.*, Sec. B, 15 (1) : 1-42.
- PETERS, W. C. H. 1854. *Naturwissenschaftliche Reise nach Mosambique in 1842 bis 1848 ausgeführt. Monatsb. Akad. Wiss. Berlin*, pp. 610-614.
- PETTIBONE, M. M. H. 1966. Revision of the Pilargidae (Annelida : Polychaeta) including descriptions of new species, and redescription of *Podarmus ploa* Chamberlin (Polynoidae). *Proc. U.S. Nat. Mus.*, 118 : 155-207.
- . 1969. Review of some species referred to *Scalissetos* McIntosh (Polychaeta Polynoidae). *Proc. Biol. Soc. Wash.*, 82 : 1-30.
- PFLUGFELDER, O. 1932. Beschreibung einiger neuer Acoetinae. Mit einem Anhang ueber eigenartige epitheliale Sinnesorgane dieser Formen. *Zool. Anz. Leipzig*, 98 : 281-295.
- . 1933. Landpolychaeten aus Niederländische-Indien. Ergebnisse der Sunda-Expedition der Notgemeinschaft der deutschen Wissenschaft 1929-30. *Zool. Anz. Leipzig*, 105 (3/4) : 65-76.
- PILLAI, T. G. 1958. Studies on a brackish water polychaetous annelid, *Marphysa borradalei*, sp. n. from Ceylon. *Ceylon J. Sci.*, 1 (2) : 94-106.
- . 1960. Some marine and brackish water serpulid Polychaeta from Ceylon, including new genera and species. *Ibid.*, 3 (1) : 1-40.
- . 1961. Annelida polychaeta of Tambalagam Lake, Ceylon. *Ibid.*, 4 (1) : 1-40.
- . 1965. Annelida polychaeta from the Philippines and Indonesia. *Ibid.*, Biol., Sec., 5 (2) : 110-177.
- . 1970. Studies on a collection of spirorbids from Ceylon, together with a critical review and revision of spirorbid systematics, and an account of their phylogeny and zoogeography. *Ibid.*, 8 (2) : 100-172.
- PLATE, L. 1916. Fauna ceylanica. Untersuchungen zur Fauna Ceylon's nach den Sammlungen von L. Plate. *Jena Zeits. Naturw.*, 54 : 1-42.
- PRASAD, M. A. S. 1969. Instances of the pelagic fish *Ilisha indica* (Swms.) feeding on the sedentary polychaete, *Prionospio pinnata* Ehlers. *J. mar. biol. Ass. India*, 9 (2) : 441-442.
- RAJA, B. T. A. 1963. Observations on the rate of growth, sexual maturity and brain of four sedentary organisms from the Madras harbour. *J. mar. biol. Ass. India*, 5 : 113-132.
- RANGARAJAN, K. 1963. On the occurrence of *Lepidasthenia ohshima* Okuda (fam. Polynoidae, Polychaeta) in Palk Bay, South India. *Ibid.*, 5 (1) : 103-107.
- . 1963. A new species of *Scyphoproctus* Gravier (Family Capitellidae) from the Gulf of Mannar, South India. *Ibid.*, 5 : 251-257.
- . 1964. A new polychaete of the family Pilargidae from Palk Bay, South India. *Ibid.*, 6 : 122-127.
- AND S. MAHADAVAN. 1961. On a new species of *Nothria* Malmgren (Polychaeta, Annelida) from the Gulf of Mannar. *Ibid.*, 3 : 179-185.

- AND C. SANKARANKUTTY 1961. A note on an abnormal *Leonnates jousseaumei* Gravier (Family Nereidae, Polychaeta). *Ibid.*, 3 : 265-266.
- RAO, K. P. 1963. Some biochemical mechanisms of low temperature acclimation in tropical poikilotherms. Abstract. *Intern. Sym. Cyto-ecol. Moscow*, pp. 50-51.
- RAO, G. C. AND P. N. GANAPATI 1966. On a new species of *Sphaerosyllis* (Polychaeta) from the beach sands of Waltair Coast. *Proc. Indian Acad. Sci.*, B. 63 (6) : 309-312.
- AND — 1967. On some interstitial polychaetes from the beach sands of Waltair Coast. *Ibid.*, 55 (B, no. 1), 10-15.
- AND — 1968. On some archiannelids from the beach sands of Waltair Coast. *Ibid.*, 67 : 24-29.
- REMSCHIED, E. 1918. Beitrage zur Kenntnis der Myzostomiden. *Abh. Senckenb. naturf. Ges. Frankfurt-am-Main*, 35 : 179-226.
- RULLIER, F. 1960. Sur deux *Ancistrosyllis groenlandica* McIntosh recueillis en Inde. *Bull. Soc. Zool. France*, 85 : 236-240.
- RUNGANATHAN. 1943. On the occurrence of hermaphroditism in *Sabellastarte magnifica*, together with an account of the excretory organs in the species. *Proc. Indian Sci. Congr.*, 29 (3) : 150-151.
- 1943. A review of the mode of breeding and development in brackish and freshwater polychaetes. *Proc. Indian Sci. Congr.*, 30 (3) : 72.
- SANDERS, H. L. 1969. Marine benthic diversity : a comparative study. *Amer. Nat.*, 102 : 243-282.
- SERRES, M. de 1855. Note sur un nouveau genre d'Annélide tubicole perforant. *Ann. Sci. Nat. Paris*, (sér. 4) 4 : 238-243.
- SILVA, P. H. D. H. de 1961. Contributions to the knowledge of the Polychaete fauna of Ceylon. Five new species, two new varieties and several new records principally from the southern coast. *Spolia Zeylanica*. *Bull. Nat. Mus. Ceylon*, 29 (2) : 164-194.
- 1965. New species and records of Polychaeta from Ceylon. *Proc. Zool. Soc. Lond.*, 144 (4) : 537-563.
- 1965. Notes on some polychaetes from Ceylon. *Spolia Zeylanica*, 30 (2) : 205-226.
- 1967. Studies on the biology of Spirorbinae (Polychaeta). *J. Zool. Proc. Zool. Soc. Lond.*, 52 (3) : 269-279.
- STEPHENSON, J. 1908. Preliminary description of an oligochaete worm of uncertain position. *Rec. Indian Mus. Calcutta*, 2 : 39.
- 1910. *Matla bengalensis* : a correction. *Ibid.*, 5 : 82. (refers to *Capitella*).
- STERZINGER, I. 1909. Einige neue *Spirorbis* Arten aus Suez. *Sitzber. Akad. Wiss. Vienna*, 118 : 1141-1159.
- STORCH, V. 1966. Drei neue Polychaeten aus dem Bitoral des Roten Meeres. *Kieler Meeresforsch.*, 22 (Heft 2) : 71-175.
- 1967. Neue Polychaeten aus der Sandfauna des Rothen Meeres. *Zool. Anz. Leipzig*, 178 (Heft 1/2) : 102-110.
- 1967. *Iphione muricata* (Savigny), ein den Chitonen aehnlicher Lebensformtyp unter den Polychaeten. *Kieler Meeresforsch.*, 23 (2) : 148-155.
- STORCH, V. AND R. NIGGEMANN 1967. Auf Echinodermen lebende Polychaeten. *Ibid.*, 23 (2) : 156-164.

- SUBRAMANIAM, M. K. 1938. On *Myzostoma gopalai* n. sp. from the Madras Harbour. *Proc. Indian Acad. Sci., Sec. B*, 7 : 270-276.
- AND R. G. AIYAR 1936. On the possible effect of the environment on the cytoplasmic inclusions in the oocytes and oögonia of *Dasychone cingulata*, *Salmacis bicolor* and *Clibanarius olivaceus*. *Ibid.*, (B), 3 : 175-194.
- TAMPI, P. R. S. 1947. The eyes of polychaetes. *Proc. 33rd Indian Sci. Congr. Bangalore*, p. 121.
- 1949. Tube-building organs of Polychaetes. *Proc. Indian Acad. Sci.*, 24 (sec. B), (4) : 129-147.
- 1958. Pelagic swarming of *Polyophthalmus* (Family Opheliidae-Polychaeta). *J. Bombay Nat. Hist. Soc.*, 55 (2) : 371-374.
- 1959. The anatomy of *Armandia leptocirris* Grube (Polychaeta). *J. Zool. Soc. Lond.*, 10 (1) : 15-32.
- 1960. On the early development of *Protula tubularia* (Montagu). *J. mar. biol. Ass. India*, 5 (1) : 53-56.
- 1963. On the swarming of heteronereids in the Gulf of Mannar. *Ibid.*, 5 (2) : 246-250.
- AND K. RANGARAJAN 1963. On the occurrence of *Arenicola brasiliensis* Nonato (Family Arenicolidae, Polychaet) in Indian waters. *Ibid.*, 5 (1) : 108-112.
- AND — 1964. Some polychaetous annelids from the Andaman waters. *Ibid.*, 6 (1) : 98-121.
- TERIO, B. 1953. Raccolte planctoniche eseguite dalla R. N. *Cherso* in Somalia. Particolare riguardo all'ermafroditismo riscontrato nella specie *Tomopteris (Johnstonella) mariana* Greeff. *Boll. Zool. Torino*, 20 : 103-106.
- THOMAS, P. J. 1963. Polychaetous worms from the Arabian Sea. A new species of the genus *Loandalia* Monro. *Bull. Dept. Mar. Biol. and Oceanogr. Univ. Kerala*, 1 : 29-34.
- 1964. *Cossura coasta* Kitamori (Polychaet : Cossuridae) : A new record from Indian waters. *Curr. Sci.*, 33 (24) : 754-755.
- TREADWELL, A. L. 1929. New species of polychaetous annelids in the collections of the American Museum of Natural History, from Porto Rico, Florida, Lower California and British Somaliland. *Amer. Mus. Novitat. N.Y.*, 392 : 1-13.
- 1943. Biological Results of the last cruise of the *Carnegie*. *Sci. Res.*, pp. 30-59.
- VAILLANT, L. 1865. Note sur un cas nouveau de reproduction par bourgeonnement, observe sur un Annélide de la rade de Suez. *C. R. Acad. Sci. Paris*, 60 : 441-443.
- WESENBERG-LUND, E. 1949. Polychaetes of the Iranian Gulf. *Danish Sci. Invest. Iran*, (4) : 247-400.
- 1958. Studies on the fauna of Curacao and other Caribbean Islands. No. 30. Lesser Antillean polychaetes, chiefly from brackish water, with a Survey and a Bibliography of Fresh and Brackish water Polychaetes. *Natuurw. Stud. Suriname*, 17 : 1-41.
- WHEELER, W. M. 1897. The sexual phases of *Myzostoma*. *Mitt. Zool. Sta. Neapel*, 12 : 228-302.
- WILLEY, A. 1906. A harbour worm and a boxing crab. *Spolia Ceylanica, Colombo*, 3 : 222-226.
- 1907. Notes on phosphorescence in marine animals, by R. E. Lloyd, with a description of a new polychaete worm. *Rec. Indian Mus. Calcutta*, 1 : 257-260.
- 1908. The fauna of brackish ponds at Port Canning, Lower Bengal. Description of a new species of polychaete worm of the genus *Splo*. *Ibid.*, 2 : 389-390.